UNIFORM A PRIORI ESTIMATES FOR A CLASS OF HORIZONTAL MINIMAL EQUATIONS

RICARDO SA EARP

ABSTRACT. In the product space $\mathbb{H}^n \times \mathbb{R}$, we obtain uniform a priori C^0 horizontal length estimates, uniform a priori C^1 boundary gradient estimates, as well as uniform modulus of continuity, for a class of horizontal minimal equations. In two independent variables, we derive an uniform global a priori C^1 estimates and we infer an existence result.

1. INTRODUCTION

In this paper, we derive uniform a priori horizontal length estimates and uniform a priori boundary gradient estimates for positive smooth solutions of a wider class of quasilinear elliptic equations, indexed by the parameter $\epsilon \in [0, 1]$. We call such equations the ϵ -horizontal minimal equations (see equation (5)). If $\epsilon = 0$, we find the horizontal minimal equation [42, Equation 32].

The uniform C^0 estimates for positive smooth solutions on bounded domains, continuous up to the boundary, is obtained by comparing, by maximum principle, such solutions with certain geometric subsolutions and supersolutions. These a priori C^0 estimates depends on the width of the domain and on the boundary value data.

In the same sprit as in [44], we obtain uniform a priori boundary gradient estimates on arbitrary bounded smooth convex domains for solutions of equation (5), that are smooth and positive up to the boundary. Indeed, we provide uniform analytic *barriers* on bounded C^0 convex domains to ensure uniform modulus of continuity for positive continuous solutions of (5) that are continuous and positive up to the boundary.

In the case of two variables, we are able to derive uniform a priori global C^1 estimates on bounded smooth convex domains, making an additional (strong) assumption on the horizontal length. However, this assumption is compatible with the geometry of the ambient space, *i.e.*, it is invariant by hyperbolic translations.

Date: April 1, 2016.

²⁰⁰⁰ Mathematics Subject Classification. 53C42, 35J25.

The author wish to thank CNPq, FAPERJ of Brazil, for a financial support.

We obtain an existence result for the ϵ -horizontal minimal equation in two independent variables on bounded smooth convex domains, taking certain smooth positive boundary value data. This existence result is, in certain sense, a counterpart of a non-existence results discussed below.

Particularly, we derive the following consequence. Given a smooth positive function f on $\partial\Omega$; if c_0 is a constant large enough, then there exists a solution of the ϵ - horizontal minimal equation in Ω , taking boundary value data $f + c_0$ on $\partial\Omega$. Actually, it suffices to take c_0 greater than the sum of the oscillation of the boundary data f with the half of the horizontal width of the domain Ω .

The techniques use suitable barriers and the maximum principle, combined with our a priori uniform C^1 estimates. Then, we are able to apply the Leray-Schaulder theorem and method [2], [17], [18] to accomplish the proof of our existence result.

Existence results for the Dirichlet problem for the ϵ -horizontal minimal equation in $n \ge 3$ variables is an open problem.

We point out that if $\epsilon > 0$, the equation is strictly elliptic, but if $\epsilon = 0$ this fact is no longer true in general. Indeed, there are many examples that can be constructed to show lack of strictly ellipticity for the horizontal minimal equation, even on bounded domains. We give now a significant example of this phenomenon in two variables. Just take $g(x,t) = x \sinh t$, 0 < x < 1, 0 < t < 1. It is easy to verify that y = g(x,t) satisfies the horizontal minimal equation. A simple verification shows that if $x \to 0$, then the first eigenvalue of the horizontal minimal operator goes to zero. On the other hand, it is amazing that the vertical minimal equation is strictly elliptic for all values of the independent variables [42, Equation 4], [45, Equation 6].

Notice that there is a non-existence result that follows from the asymptotic principle proved by E. Toubiana and the author in [45, Theorem 2.1]. Namely, there is no horizontal minimal graph given by a function $g \in C^2(\Omega) \cap C^0(\overline{\Omega})$ on a bounded smooth strictly convex domain Ω , taking zero boundary data on $\partial\Omega$. Furthermore, there is no horizontal minimal graph in $\mathbb{H}^2 \times \mathbb{R}$, over a bounded Jordan domain Ω strictly contained in an horizontal slab of height π , given by a function $g \in C^2(\Omega) \cap C^0(\overline{\Omega})$, taking zero asymptotic boundary data on $\partial\Omega$ [45, Corollary 2.1].

The theory of minimal and constant mean curvature surfaces in the product space $\mathbb{H}^2 \times \mathbb{R}$, where \mathbb{H}^2 is the hyperbolic plane is now a rich field of intense research. The notion of vertical graph has a major importance in this theory. The first main results about minimal vertical

 $\mathbf{2}$

graphs were derived by B. Nelli and H. Rosenberg [36]. Since then, a significant progress in the theory has been achieved, see, for instance, [45], [28], [27], [29]. When the ambient space is $\mathbb{H}^n \times \mathbb{R}$, the notion of vertical mean curvature equation in n independent variables has also been established and developed [49], [47], [25], [4].

On the other hand, there exists a notion of graph called "horizontal graph" that has been focused in the theory of minimal and constant mean curvature surfaces [42], [43].

The notion of horizontal graph arises naturally in the theory of hypersurfaces in $\mathbb{H}^n \times \mathbb{R}$. There are many interesting examples of minimal and constant mean curvature horizontal graphs in $\mathbb{H}^n \times \mathbb{R}$ given by explicit formulas. In fact, there are complete horizontal minimal graphs and there are entire horizontal constant mean curvature graphs [6], [7], [13], [14].

We notice, that another notion of *horizontal graph* appears naturally in the theory, as well. More precisely, they are *horizontal graph with respect to a geodesic of* \mathbb{H}^2 . This notion arises in the proof of a Schoen type result for minimal surfaces in $\mathbb{H}^2 \times \mathbb{R}$ [21].

We choose the upper half-plane model of hyperbolic plane $\mathbb{H}^2 = \{(x,y), y > 0\}$, endowed with the hyperbolic metric $d\sigma^2 = \frac{dx^2 + dy^2}{y^2}$. A horizontal graph in $\mathbb{H}^2 \times \mathbb{R}$ is the set $S = \{(x, g(x, t), t), (x, t) \in \Omega\} \subset \mathbb{H}^2 \times \mathbb{R}$, where $\Omega \subset \partial_{\infty} \mathbb{H}^2 \times \mathbb{R}$ is a domain and g(x, t) > 0, for every $(x, t) \in \Omega$. This means that in any slice of $\mathbb{H}^2 \times \mathbb{R}$ given by $t = \operatorname{cst}$, each horizontal geodesic $x = \operatorname{cst}, y > 0$ intersects S in one point at most. We call the positive function g(x, t) the horizontal length of the graph. If S is a horizontal minimal graph in $\mathbb{H}^2 \times \mathbb{R}$ the positive function g(x, t) satisfies equation (4).

It is worthwhile to notice that the horizontal minimal equation (4) is not invariant by Euclidean translations along the *y*-axis. Of course, the structure of the horizontal minimal equation does not ensure the uniqueness of the solution of the Dirichlet problem on bounded domains. Now, looking to the model minimal surfaces described in [45, Proposition 2.1] and used in [43, Proof of Theorem 1.1] to prove a Bernstein type theorem, we see that there is a family of horizontal minimal graphs over rectangles \mathcal{R} of *t*-height greater than π taking zero boundary data over $\partial \mathcal{R}$. Furthermore, over domains of arbitrarily large *x*- width, each element of this family takes arbitrarily small positive constant boundary data over a bounded domain in a rectangle \mathcal{R} , but the family attains arbitrarily large horizontal length. Thus, for the horizontal minimal equation, it follows that the dependence of the a

priori horizontal length estimates on the width of the domain is quite natural.

However, over certain admissible convex domains such that the boundary data has an admissible bounded slope condition, the solution is the Morrey's solution of the Plateau problem [30]. This result follows from the uniqueness theorem established in [43]. The uniqueness or the non-uniqueness of the Dirichlet problem for the ϵ - horizontal minimal equation on bounded convex domains and positive boundary data is an open problem.

We remark that we need to consider the family of ϵ - horizontal minimal equations in order to prove the existence of our Dirichlet problem for the horizontal minimal equation. The scheme of our construction is the following. First, we prove the existence result in the strictly elliptic situation, *i. e.* when $\epsilon > 0$. Then, we are able to deduce it for $\epsilon = 0$; that is, we get a solution for the horizontal minimal equation. In fact, in view of our uniform a priori C^1 global estimates and elliptic theory, this solution is obtained as the limit in the C^2 -topology of a sequence g_{ϵ_n} , as $0 < \epsilon_n \to 0$, satisfying the ϵ_n - horizontal minimal equation.

ACKNOWLEDGMENTS: The author warmly thanks to Eric Toubiana and Barbara Nelli for their valuable observations.

2. The ϵ -horizontal minimal equation in $\mathbb{H}^n \times \mathbb{R}$.

In this section we give some computations, some model subsolutions and supersolutions and we write down some basic properties of the ϵ -horizontal minimal equation in the product space $\mathbb{H}^n \times \mathbb{R}$.

We choose the upper half-plane model of the hyperbolic plane $\mathbb{H}^n = \{(x_1, x_2, \ldots, x_{n-1}, y), y > 0\}$, endowed with the hyperbolic metric $d\sigma^2 = \frac{dx_1^2 + \cdots + dx_{n-1}^2 + dy^2}{y^2}$. The asymptotic boundary of \mathbb{H}^n , denoted by $\partial_{\infty}\mathbb{H}^n$ is the set $\partial_{\infty}\mathbb{H}^n = \{(x_1, x_2, \ldots, x_{n-1}, y); y = 0\} \cup \{\infty\}$, see [46, Chapter 3]. The space $\mathbb{H}^n \times \mathbb{R}$ is endowed with the product metric $d\sigma^2 + dt^2$. A horizontal graph in $\mathbb{H}^n \times \mathbb{R}$ is the set

 $S = \{(x_1, x_2, \dots, x_{n-1}, g(x_1, x_2, \dots, x_{n-1}, t), t), (x_1, x_2, \dots, x_{n-1}, t) \in \Omega\} \subset \mathbb{H}^n \times \mathbb{R}, \text{ where } \Omega \subset \partial_{\infty} \mathbb{H}^n \times \mathbb{R} \text{ is a domain and}$ $g(x_1, x_2, \dots, x_{n-1}, t) > 0, (x_1, x_2, \dots, x_{n-1}, t) \in \Omega. \text{ This means that}$

at any slice of $\mathbb{H}^n \times \mathbb{R}$ given by $t = \operatorname{cst}$, each horizontal geodesic $(x_1, x_2, \ldots, x_{n-1}) = \operatorname{cst}, y > 0$ intersects S at most on one point.

We give now the computations of some important geometric quantities of the horizontal graph S. Let us consider the natural global adapted tangent field on S given by $X_{x_1} = (1, 0, \ldots, 0, g_{x_1}, 0), \ldots$,

 $X_{n-1} = (0, \ldots, 0, 1, g_{x_{n-1}}, 0), X_t = (0, \ldots, 0, g_t, 1)$. We denote by \mathfrak{g}_{ij} and by b_{ij} the coefficients of the first and second fundamental forms, respectively (with respect to the adapted frame field).

Let us define the quantity $W := g^2 \left(1 + g_{x_1}^2 + \dots + g_{x_{n-1}}^2 \right) + g_t^2.$

• The coefficients of the first fundamental form are

$$\begin{aligned} \mathbf{g}_{kk} &= \frac{1 + g_{x_k}^2}{g^2}, \ 1 \leqslant k < n, \ \mathbf{g}_{jk} = \frac{g_{x_j} g_{x_k}}{g^2}, \ 1 \leqslant j < k < n, \\ \mathbf{g}_{nk} &= \frac{g_{x_k} g_t}{g^2}, \ 1 \leqslant k < n, \ \mathbf{g}_{nn} = \frac{g^2 + g_t^2}{g^2}. \end{aligned}$$

• The inverse of the matrix (\mathfrak{g}_{ij}) is the matrix (\mathfrak{g}^{ij}) given by

$$\begin{split} \mathbf{g}^{kk} &= \left[g^2 \left(1 + g_{x_1}^2 + \dots + \widehat{g_{x_k}^2} + \dots g_{x_{n-1}}^2 \right) + g_t^2 \right] \cdot \frac{g^2}{W} \,, \\ 1 &\leqslant k < n. \\ \mathbf{g}^{kj} &= -g^2 g_{x_j} g_{x_k} \cdot \frac{g^2}{W}, \, 1 \leqslant j < k < n. \\ \mathbf{g}^{nk} &= -g_t g_{x_k} \cdot \frac{g^2}{W}, \, 1 \leqslant k < n. \\ \mathbf{g}^{nn} &= \left(1 + g_{x_1}^2 + \dots + g_{x_{n-1}}^2 \right) \cdot \frac{g^2}{W}. \end{split}$$

Let us assume that our horizontal graph S is oriented by the unit normal N given by

(1)
$$N = \left(-g_{x_1}g^2, -g_{x_2}g^2, \dots, -g_{x_{n-1}}g^2, g^2, -g_t\right) \cdot \frac{1}{W^{1/2}}$$

• The coefficients of the second fundamental form are given by

$$b_{kk} = \left(\frac{1}{g} + g_{x_k x_k} + \frac{g_{x_k}^2}{g}\right) \cdot \frac{1}{W^{1/2}}, 1 \leq k < n.$$

$$b_{kj} = \left(g_{x_k x_j} + \frac{g_{x_k} g_{x_j}}{g}\right) \cdot \frac{1}{W^{1/2}}, 1 \leq j < k < n.$$

$$b_{nk} = g_{tx_k} \cdot \frac{1}{W^{1/2}}, 1 \leq k < n.$$

$$b_{nn} = \left(g_{tt} - \frac{g_t^2}{g}\right) \cdot \frac{1}{W^{1/2}}.$$

PROPOSITION 2.1. Let S be a horizontal graph oriented by the unit normal N given by (1). Let H = H(N) be the mean curvature of S. Then the horizontal mean curvature equation is given by

(2)
$$\mathcal{M}_{\mathcal{H}}(g) = \frac{nH}{g^2} \left(g^2 \left(1 + g_{x_1}^2 + \dots + g_{x_{n-1}}^2 \right) + g_t^2 \right)^{3/2}$$

where

$$\mathcal{M}_{\mathcal{H}}(g) := \sum_{k=1}^{n-1} g_{x_k x_k} \left[g^2 \left(1 + g_{x_1}^2 + \dots + \widehat{g_{x_k}^2} + \dots + \widehat{g_{x_k}^2} + \dots + g_{x_{n-1}}^2 \right) + g_t^2 \right] + \left(1 + \sum_{k=1}^{n-1} g_{x_k}^2 \right) g_{tt}$$
$$-2 \sum_{k=1}^{n-1} g_{x_k} g_t g_{x_k t} - 2g^2 \sum_{1 \le j < k \le n-1} g_{x_j} g_{x_k} g_{x_j x_k} + (n-1)g \left(1 + \sum_{k=1}^{n-1} g_{x_k}^2 \right) + (n-2)\frac{g_t^2}{g}$$

Now, if S is a horizontal minimal graph in $\mathbb{H}^n \times \mathbb{R}$, *i.e.* H = 0, then the positive function $y = g(x_1, x_2, \dots, x_{n-1}, t)$ that we call the *horizontal length* satisfies the equation

$$(3) \quad \mathcal{M}_{\mathcal{H}}(g) = 0$$

If n = 2 we recover the horizontal minimal equation in $\mathbb{H}^2 \times \mathbb{R}$ [42, equation (2)]:

(4)
$$\mathcal{M}_{\mathcal{H}}(g) = g_{xx}(g^2 + g_t^2) + g_{tt}(1 + g_x^2) - 2g_xg_tg_{xt} + g(1 + g_x^2) = 0.$$

From now on, we focus on a 1- parameter family of elliptic equations including equation (3). Given a constant $\epsilon \in [0, 1]$, we say that a positive C^2 function g on a domain Ω is a solution of the ϵ -horizontal minimal equation, if it satisfies the following quasilinear elliptic P.D.E:

$$\mathcal{M}_{\mathcal{H}\epsilon}(g) := \sum_{k=1}^{n-1} \left[g^2 \left(1 + g_{x_1}^2 + \dots + \widehat{g_{x_k}^2} + \dots + g_{x_{n-1}}^2 \right) + g_t^2 + \frac{\epsilon}{n-1} \right] g_{x_k x_k} \\ + \left(1 + \sum_{k=1}^{n-1} g_{x_k}^2 \right) g_{tt} - 2 \sum_{k=1}^{n-1} g_{x_k} g_t g_{x_k t} - 2g^2 \sum_{1 \le j < k \le n-1} g_{x_j} g_{x_k} g_{x_j x_k} \\ + (n-1)g \left(1 + \sum_{k=1}^{n-1} g_{x_k}^2 \right) + (n-2) \frac{g_t^2}{g} \\ = 0$$

Setting
$$a_{kk}(g, Dg) := \left[g^2 \left(1 + g_{x_1}^2 + \dots + \widehat{g_{x_k}^2} + \dots + g_{x_{n-1}}^2\right) + g_t^2 + \frac{\epsilon}{n-1}\right], k = 1, \dots, n-1, \quad a_{nn}(g, Dg) = \left(1 + \sum_{k=1}^{n-1} g_{x_k}^2\right) \text{ and } a_{kn}(g, Dg) = -g_{x_k}g_t, k = 1, \dots, n-1, \quad a_{jk}(g, Dg) = -g^2 \sum_{1 \le j < k \le n-1} g_{x_j}g_{x_k}, \text{ then the symmetric matrix } a_{ij}(g, Dg), i, j = 1, \dots, n \text{ is positive, and satisfies}$$

$$(6) \sum_{i,j} a_{ij}(g, Dg)\xi_i\xi_j \leq \operatorname{trace}(a_{ij}(g, Dg))\sum_{k=1}^n \xi_k^2 = \left(1 + \epsilon + g^2(n-1) + (g^2(n-2) + 1)(g_{x_1}^2 + \dots + \dots + g_{x_{n-1}}^2) + (n-1)g_t^2\right)|\xi|^2 \leq \left(2 + g^2(n-1) + \left(\max\{1, g^2\}(n-2) + 1\right)\left(g_{x_1}^2 + \dots + g_{x_{n-1}}^2 + g_t^2\right)\right)|\xi|^2$$

And

(7)

$$\sum_{i,j} a_{ij}(g, Dg)\xi_i\xi_j = \sum_{k=1}^{n-1} \xi_k^2 (g^2 + \frac{\epsilon}{n-1}) + \xi_n^2 + g^2 \sum_{1 \le j < k \le n-1} \left(\xi_k g_{x_j} - \xi_j g_{x_k}\right)^2 + \sum_{k=1}^{n-1} \left(\xi_k g_t - \xi_n g_{x_k}\right)^2 \ge \min\{1, g^2\} |\xi|^2,$$
where $\xi \in \mathbb{R}^n \setminus \{0\}, |\xi|^2 = \sum_{k=1}^n \xi_k^2.$

By invoking inequality (7), we deduce that the ϵ -horizontal minimal equation (5) is a second order quasilinear elliptic equation. Of course, if $\epsilon > 0$, equation (5) is strictly elliptic. If $\epsilon = 0$ and if g is a C^2 positive function on a bounded domain Ω , continuous up to the boundary, then (5) is a strictly elliptic equation (cf. Section 3, inequality (8)).

REMARK 2.1. It is worth noticing that L. Hauswirth, H. Rosenberg and J. Spruck proved a half-space theorem for properly embedded constant constant mean curvature 1/2 surfaces in $\mathbb{H}^2 \times \mathbb{R}$, where the construction of H = 1/2 horizontal graphs play a significant role in the proof [20, Theorem 1.1].

We recall now the definition of subsolution and supersolution [45], [47]. We say that a C^2 function $u: \Omega \to \mathbb{R}$ is a subsolution of equation (5) in a domain Ω , if $\mathcal{M}_{\mathcal{H}_{\epsilon}}(u) \ge 0$, in Ω . We say that $w: \Omega \to \mathbb{R}$ is a supersolution if $\mathcal{M}_{\mathcal{H}_{\epsilon}}(w) \le 0$. It is well-known that elliptic theory ensures the following [39],[17], [40]:

PROPOSITION 2.2 (*Classical maximum principle*). Let $u : \Omega \to \mathbb{R}$ and $w : \Omega \to \mathbb{R}$ be a subsolution and a supersolution, for equation (5), respectively. Then if $u \leq w$ in Ω and if there is a point $p \in \Omega$ such that u(p) = w(p), it follows that u = w in Ω .

Remark 2.2.

(1) Is is easy to see that the Euclidean *n*-planes in $\mathbb{H}^n \times \mathbb{R}$ given by $y = \sum_{k=1}^{n-1} a_k x_k + bt + c; \ a, b, c \in \mathbb{R}, \ y > 0$ are positive subsolutions of equation (5). In particular the *horocylinders* given by $y = c, \ c > 0$, are subsolutions of (5).

- (2) Of course, the vertical geodesic n-planes $\mathcal{P} = \mathcal{P}(R, x_0) = \{(x_1, \dots, x_{n-1}, y, t) \in \mathbb{H}^n \times \mathbb{R}; y = \sqrt{R^2 (x_1 a_1)^2 \dots (x_{n-1} a_{n-1})^2}, t \in (-\infty, \infty), R^2 (x_1 a_1)^2 \dots (x_{n-1} a_{n-1})^2 > 0\},$ where R > 0 and $a_1, \dots, a_{n-1} \in \mathbb{R}$, are solutions of equation (3). Moreover, it is routine to check that they are supersolutions for equation (5), as well.
- (3) Minimality is invariant by a positive isometry of $\mathbb{H}^n \times \mathbb{R}$ given by a hyperbolic translation of $\mathbb{H}^n \times \{0\}$ along a horizontal geodesic L [46]. Particularly, the homothety in \mathbb{H}^n with center $(a_1, \ldots, a_{n-1}) \in \mathbb{R}^{n-1}$, ratio $\lambda > 0$, keeping \mathbb{H}^n invariant gives rise to an isometry of the product space $\mathbb{H}^n \times \mathbb{R}$ given by $(x_1, \ldots, x_{n-1}, y, t) \mapsto \left(\lambda \left[(x_1, \ldots, x_{n-1}, y) - (a_1, \ldots, a_{n-1}, 0) \right] + (a_1, \ldots, a_{n-1}, 0), t \right)$. In view of this observation, we deduce that if $y = g(x_1, \ldots, x_{n-1}, t)$ is a positive solution of (3) on a domain Ω then $y = \lambda g(\frac{x_1}{\lambda}, \ldots, \frac{x_{n-1}}{\lambda}, t), \lambda > 0$ is also a solution of (3) on $T_\lambda(\Omega)$, where T_λ is the linear map given by the matrix $\begin{pmatrix} \lambda I_{n-1} & 0 \\ 0 & 1 \end{pmatrix}$, and where I_{n-1} is the $n-1 \times n-1$ identity matrix.
- (4) Important model minimal hypersurfaces in $\mathbb{H}^n \times \mathbb{R}$ are the hypersurfaces invariant by hyperbolic translations. They are complete horizontal minimal graphs, constructed by P. Bérard and the author, if $n \ge 3$ [6, Theorem 3.8]. The minimal hypersurfaces in $\mathbb{H}^n \times \mathbb{R}$ invariant by parabolic transations are complete horizontal graphs. They were obtained by M. F. Elbert and the author [13, Theorem 3.2], if $n \ge 3$.

3. Uniform horizontal length estimates

In this section we infer uniform horizontal length estimates for solutions of equation (5) over bounded domains that are continuous and positive up to the boundary. In order to do that, we find minimal and constant mean curvature surfaces that will be used as barriers.

We need to define some geometric quantities that will be useful in the sequel.

DEFINITION 3.1. Let $\Omega \subset \partial_{\infty} \mathbb{H}^n \times \mathbb{R}$ be a bounded C^0 domain with boundary Γ . Let $f \in C^0(\Gamma)$ be a positive function. If n = 2, let

 $h(\Gamma) = \max_{\Gamma} x|_{\Gamma} - \min_{\Gamma} x|_{\Gamma}$

be the horizontal width of the domain Ω , where $x|_{\Gamma}$ is the restriction of the x coordinate to Γ . We define

$$R(\Omega, f) := \sqrt{\max_{\Gamma} f^2 + \left(\frac{h(\Gamma)}{2}\right)^2}$$

If $n \ge 3$, let Ω^{π} be the orthogonal projection of Ω on $\partial_{\infty} \mathbb{H}^n \times \{0\}$. We denote by diam (Ω^{π}) the diameter of Ω^{π} .

We define $r(\Omega)$ by the radius of the smallest n-1 closed round disk containing Ω^{π} , with center in Ω^{π} . Clearly,

 $\frac{\operatorname{diam}(\Omega^{\pi})}{2} \leqslant r(\Omega) < \operatorname{diam}(\Omega^{\pi}).$

 $\overline{W}e \ set$

$$R(\Omega, f) := \sqrt{\max_{\Gamma} f^2 + r^2(\Omega)}$$

We have therefore:

LEMMA 3.1 (Uniform horizontal length estimates).

Let $\Omega \subset \partial_{\infty} \mathbb{H}^n \times \mathbb{R}$ be a bounded C^0 domain with boundary Γ . Let $f \in C^0(\Gamma)$ be a positive function.

Assume that f admits a positive extension $g \in C^2(\Omega) \cap C^0(\overline{\Omega})$ satisfying the ϵ -horizontal minimal surface equation in Ω .

Then, the following estimate holds:

(8)
$$\min f < g < R(\Omega, f) \quad in \quad \Omega$$

Proof. If n = 2 then the proof is based on the observation that the horocylinders and the vertical geodesic planes are subsolutions and supersolutions for equation (5), respectively. Let us proceed with the proof. Let S be the graph of g. Notice that $g = y|_S$ is the horizontal coordinate y restricted to S and $f = y|_{\partial S}$ is the horizontal coordinate y restricted to ∂S .

First, we will deduce the lower bound. Let c_0 be a positive constant such that $c_0 < \min_S y|_S$. Consider the family of horocylinders $P_s, s \ge 0$ given by $y = c_0 + s, s \ge 0$. Of course, for s = 0, S is contained in the mean convex side of P_0 , that is, the horizontal length of S satisfies $y|_S > c_0$. Now letting $s \uparrow \infty$ we find a first point of contact of S with the family P_s . By the maximum principle this first point of contact should be at a point of the boundary ∂S , hence the whole surface Sshould be strictly contained in the mean convex side of the horocylinder $y = \min_{\Gamma} f$, *i.e* $y|_{S \setminus \partial S} > \min_{\Gamma} f$, or equivalently $g > \min_{\Gamma} f$ in Ω , as desired.

To obtain the upper bound, we argue as follows.

Consider the geodesic plane $\mathcal{P}(\Gamma)$ given by Remark 2.2 (2), where $x_0 =$

$$\begin{split} x_0(\Gamma) &:= \left(\max_{\Gamma} x |_{\Gamma} + \min_{\Gamma} x |_{\Gamma}\right) / 2 \text{ and } R = R(\Omega, f) = \\ \sqrt{\max_{\Gamma} (y|_{\Gamma})^2 + \left(\frac{\max x |_{\Gamma}}{2} - \frac{\min x |_{\Gamma}}{2}\right)^2}. \text{ By construction, } \partial S \subset \mathcal{P}^+(\Gamma) := \\ \{(x, y, t) \in \mathbb{H}^2 \times \mathbb{R}; \ y \leqslant \sqrt{R^2(\Omega, f) - (x - x_0(\Gamma))^2}\}. \text{ Consider now the family of geodesic planes } \mathcal{P}_s = \mathcal{P}(R(\Gamma) + s, x_0(\Gamma)), \ s \geqslant 0, \mathcal{P}_0 = \\ \mathcal{P}(\Gamma). \text{ Of course, for } s \text{ big enough, we have that } S \text{ is contained in the same connected component of } \mathbb{H}^2 \times \mathbb{R} \setminus \mathcal{P}_s(\Gamma) \text{ as } \partial S, \text{ that is } S \subset \\ \{(x, y, t) \in \mathbb{H}^2 \times \mathbb{R}; \ y < \sqrt{(R(\Gamma) + s)^2 - (x - x_0(\Gamma))^2}\}. \text{ Now letting } \\ s \downarrow 0 \text{ we cannot find a point of contact for } s > 0, \text{ by maximum principle.} \\ \text{In contrary, we should find a first interior point of contact of } S \text{ with some geodesic plane } \mathcal{P}_{s_0}. \text{ Hence, the horizontal projection of whole } S \\ \text{in } \mathbb{H}^2 \times \{0\}, \text{ would be contained in the geodesic (half Euclidean circle)} \\ y^2 + (x - x_0(\Gamma))^2 = (R(\Omega, f) + s)^2, y > 0, \text{ which gives a contradiction.} \\ \text{We conclude therefore that } S \subset \mathcal{P}^+(\Gamma). \text{ Thus, we have } y|_S \leqslant R(\Omega, f). \end{split}$$

For the same reason we cannot find an interior point of contact when, during the movement toward the original position at s = 0, the family reaches $\mathcal{P}(\Gamma)$. Indeed if this could happen then S would be a part of the geodesic plane $\mathcal{P}(\Gamma)$, so that the horizontal projection of Γ would be an arc of the geodesic $y^2 + (x - x_0(\Gamma))^2 = (R(\Omega, f))^2, y > 0$ with positive length. Since the intersection of the horizontal projection of Γ with such geodesic consists at most of two points, we get a contradiction. Henceforth, we obtain the strictly inequality $y|_{S\setminus\Gamma} < R(\Omega, f)$. This completes the proof of the Lemma, if n = 2.

Now, if n > 2 the structure of the proof is the same. Let us consider again S = graph(g). Notice that our assumption implies that ∂S lies in the side of a *n*-dimensional vertical geodesic plane $S_{R(\Omega,f)} \times \mathbb{R} \subset$ $\mathbb{H}^n \times R$, whose asymptotic boundary contains Ω^{π} . Where $S_{R(\Omega,f)}$ is a n-1 geodesic plane in \mathbb{H}^n (Euclidean n-1 halfsphere) of Euclidean radius $R(\Omega, f)$.

We now accomplish the proof, as follows: We can use the n dimensional horocylinders to obtain the horizontal lower length bounds for (5), in the same way as in the case n = 2. Moreover, the n-dimensional vertical totally geodesic planes (Remark 2.2 (2)) can be employed, to obtain the desired horizontal upper bounds for (5), working similarly as in the case n = 2.

REMARK 3.1. The proof shows that the estimate hold if we allow the solutions to be nonnegative on the boundary.

4. Uniform boundary gradient estimates and modulus of continuity

In this section we build uniform *barriers* at any point of the boundary of a bounded convex smooth domain, for a positive smooth solution g of the ϵ -horizontal minimal equation C^1 up to the boundary. We obtain in fact these a priori gradient estimates in the sprit of [44], on account of the techniques explained by Gilbarg and Trudinger [17]. We also construct analytic barriers to get an uniform *modulus of continuity* along the boundary of a bounded convex domain, for a positive solution g of the ϵ -horizontal minimal equation C^0 up to the boundary.

DEFINITION 4.1. We say that a C^0 domain $\Omega \subset \partial_{\infty} \mathbb{H}^n \times \mathbb{R}$ is convex if, for any $p \in \partial\Omega$, Ω lies in one side of some n - 1-plane Π of $\partial_{\infty} \mathbb{H}^n \times \mathbb{R}$ passing through $p, i.e \ p \in \Pi \cap \partial\Omega$ and $\Pi \cap \Omega = \emptyset$.

We need now the definition of the quantity $R(\Omega, f)$ raised in Definition 3.1.

THEOREM 4.1 (Uniform boundary gradient estimates I). Let $\Omega \subset \partial_{\infty} \mathbb{H}^n \times \mathbb{R}$ be a C^2 bounded domain. Let $\varphi \in C^2(\overline{\Omega})$ be a positive function. Let $g \in C^2(\Omega) \cap C^1(\overline{\Omega})$ be a positive solution of the ϵ -horizontal minimal equation (5) in Ω , such that $g = \varphi$ on $\Gamma = \partial \Omega$, $\epsilon \in [0, 1]$. Assume that Ω is convex. Then, the following estimate holds.

(9)
$$\max_{\Gamma} |Dg| \leqslant C$$

where
$$C = C(\min_{\Gamma} g, \max_{\Gamma} g, r(\Omega), \max_{\Omega} \varphi, \max_{\Omega} |D\varphi|, \max_{\Omega} |D^2\varphi|).$$

Proof. Let g be a solution of the ϵ -horizontal minimal equation as in the statement of the Theorem.

It suffices to get a priori estimates for the normal derivatives at any point $p \in \partial \Omega$. We obtain first the upper bound for the normal derivatives constructing an upper barrier.

We define

(10)
$$\psi(d) := \frac{1}{b_1} \ln \left(1 + \frac{e^{(R(\Omega, f) + \max_{\Omega} \varphi) b_1} - 1}{\delta_1} d \right)$$

where b_1 and δ_1 are positive constants to be defined later.

Let $p \in \partial\Omega$ and let Π be the n-1-plane passing through p as in Definition 4.1 and let $d(q) = d(q, \Pi), q \in \Omega$, where $d(q, \Pi)$ is the Euclidean distance from q to Π . We define $v(q) := \psi(d(q)), q \in \Omega$. Of course, v(p) = 0.

Let $w = \varphi + v$ and let $\overline{\mathcal{M}}_{\mathcal{H}\epsilon}$ be the strictly elliptic operator given by (11)

$$\overline{\mathcal{M}}_{\mathcal{H}_{\epsilon}}(w) := \sum_{k=1}^{n-1} \left[g^2 \left(1 + w_{x_1}^2 + \dots + \widehat{w_{x_k}^2} + \dots + w_{x_{n-1}}^2 \right) + w_t^2 + \frac{\epsilon}{n-1} \right] w_{x_k x_k} \\ + \left(1 + \sum_{k=1}^{n-1} w_{x_k}^2 \right) w_{tt} - 2 \sum_{k=1}^{n-1} w_{x_k} w_t w_{x_k t} - 2g^2 \sum_{1 \le j < k \le n-1} w_{x_j} w_{x_k} w_{x_j x_k} \\ + (n-1)g \left(1 + \sum_{k=1}^{n-1} w_{x_k}^2 \right) + (n-2) \frac{w_t^2}{g} \\ := a_{ij}(g, Dw) D_{ij} w + (n-1)g \left(1 + \sum_{k=1}^{n-1} w_{x_k}^2 \right) + (n-2) \frac{w_t^2}{g}$$
(using summation convention)

(using summation convention)

In view of inequalities (7) and (6) we have:

(12)
$$\min\{1, g^2\} |\xi|^2 \leq a_{ij}(g, Dw) \xi_i \xi_j \leq (2+g^2(n-1)+(\max\{1, g^2\}(n-2)+1) |Dw|^2) |\xi|^2 \operatorname{in} \Omega$$

Now by invoking the horizontal length estimates (Lemma 3.1), it follows that

(13)
$$\min\{1, \min_{\Gamma} g^2\} |\xi|^2 \leq a_{ij}(g, Dw) \xi_i \xi_j \quad \text{in } \Omega$$

Set $\Lambda_1 := 2 + g^2(n-1) + \left(\max\{1, g^2\}(n-2) + 1\right) |Dw|^2$
 $= 2 + g^2(n-1) + \left(\max\{1, g^2\}(n-2) + 1\right) |D\varphi + Dv|^2 \text{ in } \Omega.$

We deduce the following inequalities:

$$\begin{split} &(14) \\ \Lambda_1 < 2 + R^2(\Omega, f)(n-1) + \left(\max\{1, R^2(\Omega, f)\}(n-2) + 1 \right) |D\varphi + Dv|^2 \\ &< 2 + R^2(\Omega, f)(n-1) + 2 \left(\max\{1, R^2(\Omega, f)\}(n-2) + 1 \right) (|D\varphi|^2 + |Dv|^2) \\ &\leqslant 2 + R^2(\Omega, f)(n-1) + 2 \left(\max\{1, R^2(\Omega, f)\}(n-2) + 1 \right) |D\varphi|^2 \\ &+ 2 \left(\max\{1, R^2(\Omega, f)\}(n-2) + 1 \right) |Dv|^2 \\ &\leqslant \left(2 + R^2(\Omega, f)(n-1) + 2 \left(\max\{1, R^2(\Omega, f)\}(n-2) + 1 \right) |D\varphi|^2 \right) |Dv|^2 \\ &+ 2 \left(\max\{1, R^2(\Omega, f)\}(n-2) + 1 \right) |Dv|^2, \quad \text{whenever} \quad |Dv| \ge 1 \\ &\leqslant \left(2 + R^2(\Omega, f)(n-1) + 2 \left(\max\{1, R^2(\Omega, f)\}(n-2) + 1 \right) (|D\varphi|^2 + 1) \right) |Dv|^2, \\ &\qquad \text{whenever} \quad |Dv| \ge 1 \end{split}$$

Set
(15)
$$\alpha_1 := \frac{\left(2 + R^2(\Omega, f)(n-1) + 2\left(\max\{1, R^2(\Omega, f)\}(n-2) + 1\right)\left(\max_{\Omega} |D\varphi|^2 + 1\right)\right)}{\min\{1, \min_{\Gamma} g^2\}}.$$

By combining (13), (14) and (15) we have

(16)
$$\Lambda_1 < \alpha_1 \min\{1, \min_{\Gamma} g^2\} |Dv|^2$$

$$< \alpha_1 a_{ij}(g, D\varphi + Dv) D_i v D_j v, \quad \text{whenever} \quad |Dv| \ge 1$$

$$< \alpha_1 \mathcal{F}, \quad \text{whenever} \quad |Dv| \ge 1$$

 where
$$\mathcal{F} := a_{ij}(g, D\varphi + Dv) D_i v D_j v.$$

Now set $\Lambda_2 := 1 + |Dw|^2 = 1 + |D\varphi + Dv|^2$. In the same way as in proof of the inequalities (14), we deduce

(17)
$$\Lambda_2 < \alpha_2 \mathcal{F}$$
, whenever $|Dv| \ge 1$
where $\alpha_2 = \frac{3 + 2 \max_{\Omega} |D\varphi|^2}{\min\{1, \min_{\Gamma} g\}}$

From (11), (14), and from the definitions of w, \mathcal{F} , α_1 and α_2 , we get (18)

$$\overline{\mathcal{M}}_{\mathcal{H}\epsilon}(\varphi+v) = a_{ij}D_{ij}v + a_{ij}D_{ij}\varphi + (n-1)g\left(1 + \sum_{k=1}^{n-1}(\varphi_{x_k} + v_{x_k})^2\right) \\ + (n-2)\frac{(\varphi_t + v_t)^2}{g} \\ = a_{ij}\psi''(d)D_idD_jd + a_{ij}D_{ij}\varphi + (n-1)g\left(1 + \sum_{k=1}^{n-1}(\varphi_{x_k} + v_{x_k})^2\right) \\ + (n-2)\frac{(\varphi_t + v_t)^2}{g} \\ \text{(since d is linear)} \\ \leqslant a_{ij}\frac{\psi''(d)}{\psi'(d)^2}D_ivD_jv + \Lambda_1|D^2\varphi| + (n-1)g\left(1 + \sum_{k=1}^{n-1}(\varphi_{x_k} + v_{x_k})^2\right) \\ + (n-2)\frac{(\varphi_t + v_t)^2}{g} \\ \psi''(d) = \sum_{k=1}^{n-1}(\varphi_{x_k} + v_{x_k})^2 \right)$$

$$\leqslant a_{ij} \frac{\psi'(d)^2}{\psi'(d)^2} D_i v D_j v + \Lambda_1 |D \varphi| + (n-1)g \left(1 + \sum_{k=1}^{\infty} (\varphi_{x_k} + v_{x_k})\right)$$

$$+ (n-2) \frac{(\varphi_t + v_t)^2}{g}$$

$$< a_{ij} \frac{\psi''(d)}{\psi'(d)^2} D_i v D_j v + \Lambda_1 |D^2 \varphi|$$

$$+ \max\left\{ (n-1) \max_{\Omega} g, \frac{n-2}{\min g} \right\} (1 + |D\varphi + Dv|^2)$$

$$\leqslant -b_1 a_{ij} D_i v D_j v + \Lambda_1 |D^2 \varphi|$$

$$+ \max\left\{ (n-1) \max_{\Omega} g, \frac{n-2}{\min g} \right\} (1 + |D\varphi + Dv|^2)$$

$$< -b_1 \mathcal{F} + (\max_{\Omega} |D^2 \varphi|) \alpha_1 \mathcal{F} + \max\left\{ (n-1) \max_{\Omega} g, \frac{n-2}{\min g} \right\} \alpha_2 \mathcal{F}$$

$$< (-b_1 + \alpha) \mathcal{F}, \quad \text{whenever} \quad |Dv| \ge 1,$$

where $\alpha := (\max_{\Omega} |D^2 \varphi|) \alpha_1 + \max\left\{ (n-1) \max_{\Omega} g, \frac{n-2}{\min_{\Gamma} g} \right\} \alpha_2$

Now we choose b_1 and δ_1 such that

(1) $b_1 \ge \alpha$.

(2)
$$\frac{e^{(R(\Omega,f)+\max_{\Omega}\varphi)b_{1}}-1}{\delta_{1}} \ge b_{1}e^{(R(\Omega,f)+\max_{\Omega}\varphi)b_{1}}.$$

With these choices of b_{1} and δ_{1} we get that $|Dv| = \psi'(d) \ge 1$
if $d \le \delta_{1}.$

Define $\mathcal{N} = \{q \in \Omega; d(q) < \delta_1\}$. From the choices of b_1 and δ_1 above we deduce that $w = \varphi + v$ is a positive supersolution of (11); that is, $\overline{\mathcal{M}}_{\mathcal{H}\epsilon}(\varphi + v) < 0$ in \mathcal{N} . Observe that the linear elliptic operator \mathcal{L} given by $\mathcal{L}[g - (\varphi + v)] := \overline{\mathcal{M}}_{\mathcal{H}\epsilon}(g) - \overline{\mathcal{M}}_{\mathcal{H}\epsilon}(\varphi + v) = \mathcal{M}_{\mathcal{H}\epsilon}(g) - \overline{\mathcal{M}}_{\mathcal{H}\epsilon}(\varphi + v) = -\overline{\mathcal{M}}_{\mathcal{H}\epsilon}(\varphi + v) > 0$ in \mathcal{N} , since g is a positive solution of the ϵ -horizontal minimal equation, by assumption. Furthermore, $\mathcal{L}[g - (\varphi + v)]$ satisfies the Hopf maximum principle. Recall now that $g < R(\Omega, f)$, by Lemma 3.1. On $\partial \mathcal{N} \cap \Omega$ we get $g < R(\Omega, f) \leq \varphi +$ $R(\Omega, f) + \max \varphi = \varphi + v$. On $\partial \Omega \cap \partial \mathcal{N}$ we have $g = \varphi \leq \varphi + v$. Hence $g \leq$ $\varphi + v$ in \mathcal{N} , by the maximum principle. Moreover, $g(p) = \varphi(p) + v(p)$. Therefore, $w = \varphi + v$ is an upper barrier (since $g(q) \leq w(q), q \in \mathcal{N}$ and g(p) = w(p)). From this, the upper bound for the normal derivatives follows. We obtain henceforth the desired boundary gradient upper bound.

To obtain the lower bound for the normal derivatives, we will construct a lower barrier. Consider $\varphi - v$. Note that $\varphi(p) - v(p) = \varphi(p) = g(p) > 0$. Notice that there exists a connected part N of \mathcal{N} , containing p, such that $\varphi - v > 0$ in N. We derive the following computations.

(19)

$$\overline{\mathcal{M}}_{\mathcal{H}\epsilon}(\varphi - v) = -a_{ij}D_{ij}v + a_{ij}D_{ij}\varphi + (n-1)g\left(1 + \sum_{i=1}^{n-1}(\varphi_{x_k} - v_{x_k})^2\right) + (n-2)\frac{(\varphi_t - v_t)^2}{g}$$

$$= b_1a_{ij}D_ivD_jv + a_{ij}D_{ij}\varphi + (n-1)g\left(1 + \sum_{i=1}^{n-1}(\varphi_{x_k} - v_{x_k})^2\right) + (n-2)\frac{(\varphi_t - v_t)^2}{g}$$

 $\geq b_1 \min\{1, \min_{\Gamma} g^2\} |Dv|^2 - \max_{\Omega} |D^2 \varphi| \Lambda_1$ Thus

 $\overline{\mathcal{M}}_{\mathcal{H}_{\epsilon}}(\varphi - v) > 0, \quad \text{whenever} \quad |Dv| \ge 1,$

if b_1 is chosen big enough. So with this choice of b_1 , it follows that $\varphi - v$ is a subsolution of (11). Noticing that on $\partial \mathcal{N} \cap \Omega$, $v = \psi(\delta_1) = \max_{\Omega} \varphi + R(\Omega, f)$, we deduce that on $\partial \mathcal{N} \cap \Omega$ we have $\varphi - v = \varphi - \max_{\Omega} \varphi - R(\Omega, f) < 0 < g$. And on $\partial \Omega \cap \partial \mathcal{N}$, we have $\varphi - v \leq \varphi = g$, hence $\varphi - v \leq g$ on \mathcal{N} , by the maximum principle and $g(p) = \varphi(p) - v(p)$. We thus infer the a priori lower bound for the normal derivatives. Therefore we obtain the desired a priori boundary gradient estimates. This completes the proof of the Theorem.

REMARK 4.1. For a $C^{2,\alpha}$ domain Ω for some $0 < \alpha < 1$ whose boundary has positive mean curvature, we can find a simpler and more geometric a priori lower bounds for the gradient. In fact, these domains satisfies a *boundary slope condition* and the Euclidean *n*-planes can be used as lower barriers, since they are subsolutions of the ϵ -horizontal mean equation (5).

Recall now that the minimal equation in Euclidean space is given by

(20)
$$\operatorname{div}\left(\frac{\nabla u}{W(u)}\right) := \sum_{i=1}^{n} \frac{\partial}{\partial x_i} \left(\frac{u_{x_i}}{\sqrt{1 + \|\nabla u\|_{\mathbb{R}^n}^2}}\right) = 0$$

DEFINITION 4.2. Let $\Omega \subset \mathbb{R}^n$ be a $C^{2,\alpha}$ bounded convex domain for some $0 < \alpha < 1$, with boundary Γ . Let $f \in C^{2,\alpha}(\Gamma)$. We define by $\widehat{\varphi} \in C^{2,\alpha}(\overline{\Omega})$ the unique minimal solution of the minimal equation in Euclidean space in Ω , taking the prescribed boundary data f on Γ , given by Jenkins and Serrin [23].

We recall that the maximum principle and the use of the Euclidean *n*-planes as barriers ensures that $\min_{\Gamma} f \leq \widehat{\varphi} \leq \max_{\Gamma} f$ in Ω .

The following Lemma is well-known and we will use it in the proof of the next Lemma. We will write a proof for completeness.

LEMMA 4.1. Let $\Omega \subset \mathbb{R}^n$ be a $C^{2,\alpha}$ bounded convex domain for some $0 < \alpha < 1$. Let $\Gamma = \partial \Omega$ and let $f \in C^{2,\alpha}(\Gamma)$ be a positive function.

For each $s \in [1/2, 1]$, let $\widehat{\varphi_s} \in C^{2,\alpha}(\overline{\Omega})$ be the unique solution of (20) taking the prescribed boundary value data $(2s-1)f + 2(1-s)\min_{\Gamma} f$ on Γ . Then there exists a constant K, independent of s, such that

(21)
$$\max_{\overline{\Omega}} |D\widehat{\varphi_s}| + \max_{\overline{\Omega}} |D^2\widehat{\varphi_s}| \leqslant K$$

Proof. For the readers convenience, we outline a proof, as follows. First, note that the Extension Lemma [17, Lemma 6.38] provides a

 $C^{2,\alpha}(\overline{\Omega})$ extension φ_s of $(2s-1)f + 2(1-s)\min_{\Gamma} f$ such that $\max_{\overline{\Omega}} |\varphi_s| + \max_{\overline{\Omega}} |D\varphi_s| + \max_{\overline{\Omega}} |D^2\varphi_s|$ is bounded by a constant C independent of $s, s \in [1/2, 1]$.

As we have observed before, the maximum principle yields $\min_{\Gamma} f \leq \widehat{\varphi_s} \leq \max_{\Gamma} f$, since this inequality occurs on the boundary. Note also that that the first eigenvalue of the matrix associated to equation (20) is 1. Now using the extension φ_s , we can follow, step by step, the proof of Theorem 4.1, to ensure the bounds for the first derivatives of $\widehat{\varphi_s}$, independent of s. Then by applying the global a priori Hölder estimates of Ladyzhenskaya and Ural'tseva [24], [17, Theorem 13.7], we have global Hölder a priori estimates for the first derivatives of $\widehat{\varphi_s}$, independent of s. Finally, the global a priori Schauder estimates [17, Theorem 6.6] shows that the $C^{2,\alpha}(\overline{\Omega})$ norm $|\widehat{\varphi_s}|_{C^{2,\alpha}(\overline{\Omega})}$, is bounded by a constant K, independent of $s, s \in [1/2, 1]$. This gives the desired estimate.

Let $\ell : [0,1] \to \mathbb{R}$ be a continuous function satisfying $0 \leq \ell(s) \leq 1$, $\ell(1) = 1$ and $\ell(s) = 0$, $s \in [0, 1/2]$. For each $s \in [0, 1]$, let us now turn attention to the positive solutions of the following P.D.E:

$$(22) \quad \sum_{k=1}^{n-1} \left[g^2 \left(1 + g_{x_1}^2 + \dots + \widehat{g_{x_k}^2} + \dots g_{x_{n-1}}^2 \right) + g_t^2 + \frac{\epsilon}{n-1} \right] g_{x_k x_k} \\ + \left(1 + \sum_{k=1}^{n-1} g_{x_k}^2 \right) g_{tt} - 2 \sum_{k=1}^{n-1} g_{x_k} g_t g_{x_k t} - 2g^2 \sum_{1 \le j < k \le n-1} g_{x_j} g_{x_k} g_{x_j x_k} \\ + \left[(n-1)g \left(1 + \sum_{k=1}^{n-1} g_{x_k}^2 \right) + (n-2) \frac{g_t^2}{g} \right] \ell(s) \\ = 0$$

Note that for s = 1, equation (22) reduces to the ϵ -horizontal minimal equation.

For later purposes, we need a slight refined generalization of the boundary gradient estimates for $C^{2,\alpha}$ domains and boundary data, on account of Lemma 4.1 and the above observations.

THEOREM 4.2 (Uniform boundary gradient estimates II). Let $\Omega \subset \partial_{\infty} \mathbb{H}^n \times \mathbb{R}$ be a $C^{2,\alpha}$ bounded convex domain for some $0 < \alpha < 1$. Let $\Gamma = \partial \Omega$ and let $f \in C^{2,\alpha}(\Gamma)$ be a positive function.

For each $s \in [1/2, 1]$, let $g_s \in C^{2,\alpha}(\overline{\Omega})$ be a positive solution of (22) in Ω , such that $g_s = (2s - 1)f + 2(1 - s) \min_{\Gamma} f$ on Γ .

Then, for each $s \in [1/2, 1]$ the following estimate holds.

(23)
$$\min_{\Gamma} f \leq \max_{\overline{\Omega}} g_s < R(\Omega, f) \quad on \quad \overline{\Omega}$$

where $R(\Omega, f)$ is given by Definition 3.1 and $\max_{\Gamma} |Dg_s| \leq C$

where
$$C = C(\min_{\Gamma} f, \max_{\Gamma} f, r(\Omega), K),$$

where K is given by Lemma 4.1.

Proof. Let $\widehat{\varphi_s}$ be the unique minimal Euclidean extension of $(2s-1)f + 2(1-s) \min_{\Gamma} f$ satisfying equation (20).

Notice that, as in the proof of the Theorem 4.1, we have the following ingredients:

1: First, it is routine to check that $\min_{\Gamma} f \leq g_s|_{\Gamma} \leq \max_{\Gamma} f$. On the other hand, observe that the Euclidean *n*-planes are subsolutions of (22). Moreover, the vertical geodesic *n*-planes (Remark 2.2 (2)) are still supersolutions of (22), since $0 \leq \ell(s) \leq 1$,. Consequently, the length estimate inferred in Section 3 hold with exactly the same statement as in Lemma 3.1. Thus, by invoking the length estimate, it follows that $\min_{\Gamma} f < g_s < R(\Omega, f)$ on Ω and $\min_{\Gamma} f \leq g_s < R(\Omega, f)$ on $\overline{\Omega}$.

2: Secondly, taking into account again that $0 \leq \ell(s) \leq 1$, inequality (21), and inequalities (7) and (6), we are able to follow the procedure of the proof of Theorem 4.1 to obtain the desired estimate.

Noticing that in view of Lemma 4.1 and its proof, we have K in the place of $\max_{\Omega} |D\widehat{\varphi_s}| + \max_{\Omega} |D^2\widehat{\varphi_s}|$ and $\max_{\Gamma} f$ instead of $\max_{\Omega} \widehat{\varphi_s}$ in the estimate.

This accomplishes the proof of the theorem.

Next, we show that a positive solution $g \in C^2(\Omega) \cap C^0(\overline{\Omega})$ of (5) has a uniform modulus of continuity along a bounded convex domain Ω .

Let $\Gamma \subset \partial_{\infty} \mathbb{H}^n \times \mathbb{R}$ be a bounded convex hypersurface and let $f \in C^0(\Gamma)$. Given $\varepsilon > 0$, the continuity of f yields the existence of a positive constant $\delta_0 > 0$, such that $|f(p') - f(p)| < \varepsilon/3$, if $|p'-p| < \delta_0, p', p \in \Gamma$.

We now define the barriers φ^{\pm} at $p \in \Gamma$ that we need in the next theorem. Let

(24)
$$\varphi^{\pm}(q) := f(p) \pm \frac{\varepsilon}{3} \pm R(\Omega, f) \frac{\ln(1 + |q - p|^2)}{\ln(1 + \delta_0^2)}, \quad q \in \Omega$$

where $\delta_0 = \delta_0(f)$ is defined in the above paragraph.

THEOREM 4.3 (Uniform modulus of continuity). Let $\Omega \subset \partial_{\infty} \mathbb{H}^n \times \mathbb{R}$ be a C^0 bounded domain. Let $\Gamma = \partial \Omega$ and let $f \in C^0(\Gamma)$ be a positive function. Let $g \in C^2(\Omega) \cap C^0(\overline{\Omega})$ be a positive solution of the ϵ -horizontal minimal equation (5) in Ω , such that g = f on Γ . Assume that Ω is convex.

Then, it follows that given $\varepsilon > 0$ there is $\delta > 0$ such that if $p \in \partial \Omega$ and $q \in \Omega$, satisfy $|q - p| < \delta$, then $|g(q) - g(p)| < \varepsilon$, where $\delta = \delta(\varepsilon, \delta_0, \min_{\Gamma} g, \max_{\Gamma} g, r(\Omega), \max_{\Omega} |D\varphi^{\pm}|, \max_{\Omega} |D^2\varphi^{\pm}|).$

Proof. The proof is carry out by mimicking the proof of the boundary gradient estimates, using the barriers φ^{\pm} (instead of the function φ). Let $\varepsilon > 0$.

Notice that $f \leq \varphi^+$ on Γ and $f \geq \varphi^-$ on Γ .

We need to prove that we can find a suitable constant $\delta > 0$ such that if $|q - p| < \delta$, $q \in \Omega, p \in \Gamma$, then both inequalities $g(q) < f(p) + \varepsilon$ and $g(q) > f(p) - \varepsilon$ hold.

To obtain the first inequality we use the barrier $w^+ = \varphi^+ + v^+$, where the function v^+ is defined by $v^+(q) := \psi^+(d(q)), q \in \Omega$, where

(25)
$$\psi^+(d) := \frac{1}{b_1} \ln\left(1 + \frac{e^{R(\Omega, f)b_1} - 1}{\delta_1}d\right)$$

Indeed working exactly as in the proof of Theorem 4.1, we can deduce that w^+ is a supersolution of (11) in \mathcal{N} , and $g \leq \varphi^+ + v^+$ in \mathcal{N} , where \mathcal{N} is defined in the proof of Theorem 4.1. Next, we are able to choose $\delta > 0$ small enough, $\delta < \delta_0$, $\delta < \delta_1$, such that if $|q - p| < \delta$, then $0 \leq \varphi^+(q) < f(p) + 2\varepsilon/3$ and $0 < v^+(q) < \varepsilon/3$. We infer therefore that if $|q - p| < \delta$ then $g(q) < f(p) + \varepsilon$, as desired.

In order to achieve the proof of the theorem, we now define $v^{-}(q) := \psi^{-}(d(q)), q \in \Omega$, where

(26)
$$\psi^{-}(d) := \frac{1}{b_{1}} \ln \left(1 + \frac{e^{(R(\Omega, f) + \max_{\Omega} |\varphi^{-}|) b_{1}} - 1}{\delta_{1}} d \right)$$

Then, we can use the barrier $w^- = \varphi^- - v^-$, working as before, to deduce that w^- is a subsolution of (11) in \mathcal{N} , and finally, to obtain that $g(q) > f(p) - \varepsilon$. This accomplishes the proof of the Theorem.

5. Uniform global gradient estimates

In this section we are able to obtain uniform a priori global gradient estimates for the ϵ -horizontal minimal equation in two independent variables. The a priori interior gradient bound for the minimal equation in Euclidean space in two independent variables was established by R. Finn [15]. The *n*-dimensional case was done by Bombieri, De Giorgi and Miranda [8]. Equations of minimal type were first studied by Finn [15]. He established a priori estimates for the gradient of a solution. Then H. Jenkins and J. Serrin [22] obtained curvature estimates for some kind of such equations.

When the ambient is the hyperbolic space, R. López and S. Montiel, under certain geometric assumptions, obtained one of the first estimate of the gradient for the constant mean curvature equation. They applied it to solve a related Dirichlet problem [26].

In his fundamental paper L. Simon [48] derived a priori gradient estimates for general mean curvature type equations in two variables. His results has been applied to many mean curvature equations in two variables in several spaces [44], [12]. However, due to the geometry of the ambient space, the horizontal minimal equation does not fit the structure conditions of the equations of mean curvature type, established by L. Simon. It does not match either the structure conditions in [17, Theorem 5.2].

Moreover, the technique developed by L. Cafarelli, L. Nirenberg and J. Spruck in [9] to obtain a priori global gradient estimates, cannot be applied to the present situation. On the contrary, the method in [9] has been applied to a mean curvature equation in hyperbolic space in [33] and [5] and to a constant mean curvature equation in warped product [11]. We notice that, in some cases, the a priori global C^1 estimates depends on the derivation of both a priori C^0 estimates and a priori C^1 boundary gradient estimates, see, for instance, [32], [19] and [1].

We remark that the a priori gradient estimates inferred by J. Spruck [49], are adapted for the vertical mean curvature equation to study many problems in the product spaces $M \times \mathbb{R}$, where M is a Riemannian manifold [7], [16], [47], [10].

At last, the general a priori gradient estimates carry out by H. Rosenberg, E. Toubiana and R. Souam [41, Theorem 3.6] has been applied to solve Dirichlet problems in homogeneous spaces [37].

Next, we follow the techniques derived by Bakel'man [3]. We do the analysis in two independent variables, for the reasons that will be clear in the proof.

Notice that certain model horizontal minimal graphs in two independent variables are *Killing graphs* [45],[43]. That is they are graphs with respect to the coordinates system given by the 1-parameter group of isometries $(x, y, t) \rightarrow (\lambda x, \lambda y, t), \lambda > 0$ of $\mathbb{H}^2 \times \mathbb{R}$. This group is constituted of hyperbolic translations of the ambient space. Noticing also that by employing the translations $(\overline{x}, \overline{y}, t) \rightarrow (\lambda \overline{x}, \lambda \overline{y}, t) := (x, y, t),$ given a solution $g(\overline{x}, t)$ of the ϵ -horizontal minimal equation (27), we obtain another function $y = g_{\lambda}(x, t)$, satisfying (27), for $\epsilon \lambda^2$ in the place of ϵ (see (45)).

Due to the techniques employed, we are forced to make a strong constraint on the horizontal length to obtain the uniform global C^1 gradient estimates. Nevertheless, the assumption is invariant by hyperbolic translations, so it is compatible with the geometry of the ambient space.

We recall that if n = 2 the ϵ -horizontal minimal equation becomes

(27)
$$\mathcal{M}_{\mathcal{H}\epsilon}(g) = g_{xx}(g^2 + g_t^2 + \epsilon) + g_{tt}(1 + g_x^2) - 2g_xg_tg_{xt} + g(1 + g_x^2) = 0,$$

THEOREM 5.1 (Uniform global gradient estimates I). Let $\Omega \subset \partial_{\infty} \mathbb{H}^2 \times \mathbb{R}$ be a C^1 bounded domain. Let $g \in C^{2,\alpha}(\Omega) \cap C^1(\overline{\Omega})$ be a positive solution of the ϵ -horizontal minimal equation (27) in Ω . Assume there exist constants c_1 and c_2 , and c_3 , independent of ϵ , such that $0 < c_1 \leq g \leq c_2$ on $\overline{\Omega}$ and $|Dg| \leq c_3$ on $\partial\Omega$. Assume further that $c_2 < c_1 + c_1 \sqrt{\frac{\pi}{2}}$. Then there exists a constant C depending only on c_1, c_2 and c_3 such that $|Dg| \leq C$ on the whole $\overline{\Omega}$.

Proof. Let g be a solution of the ϵ -horizontal minimal equation on Ω , C^1 up to the boundary, such that $0 < c_1 \leq g \leq c_2$ on $\overline{\Omega}$. Assume first that $c_1 \leq 1$.

By assumption, we have $c_2 - c_1 < c_1 \sqrt{\frac{\pi}{2}}$. We first recall the elementary identity $\int_{0}^{+\infty} e^{-\gamma s^2} ds = \left(\frac{\pi}{2 \cdot 2\gamma}\right)^{1/2}$. We now consider the function $\phi(u)$ given by

(28)
$$\phi(u) = c_1 + \int_0^u e^{-\gamma s^2} \,\mathrm{d}s$$

where γ is chosen such that $c_1 > \frac{1}{\sqrt{2\gamma}} > \sqrt{\frac{2}{\pi}} \cdot (c_2 - c_1)$. It follows that $c_2 < c_1 + \left(\frac{\pi}{2 \cdot 2\gamma}\right)^{1/2}$. Let $c_{12} = c_{12}(c_1, c_2)$ be a constant such that $c_1 + \int_{-\infty}^{c_{12}} e^{-\gamma s^2} ds > c_2$.

With this choice, we are able to write

(29)
$$g(p) = \phi(u(p)), \ u \ge 0, \ p \in \overline{\Omega}$$

for some auxiliary function u satisfying $u(p) < c_{12}$ on $\overline{\Omega}$. Then, we have the inequalities

$$e^{-\gamma c_{12}^2} < \phi'(u) \leqslant 1$$

and

$$\frac{0 < -\phi''(u) \leqslant +2\gamma c_{12}}{-\phi'''(u)\phi'(u) + \phi''^{2}(u)} = 2\gamma$$

Next differentiate (29) with respect to x and t to obtain

(31)
$$g_x = \phi'(u)u_x, \ g_t = \phi'(u)u_t$$
$$g_x^2 + g_t^2 = \phi'^2(u)(u_x^2 + u_t^2)$$

Of course, with the aid of equation (31), we infer the following inequalities:

(32)
$$\sqrt{u_x^2 + u_t^2} \leqslant \frac{\max_{\partial \Omega} \sqrt{g_x^2 + g_t^2}}{\min_{\overline{\Omega}} \phi'(u)}$$
, on $\partial \Omega$

and

$$\sqrt{g_x^2 + g_t^2} \leqslant \sqrt{u_x^2 + u_t^2}, \quad \text{on} \quad \overline{\Omega}.$$

Thus, if the maximum of Du on $\overline{\Omega}$ is achieved at the boundary then $\max_{\overline{\Omega}} |Du| \leqslant \frac{\max_{\partial \Omega} |Dg|}{\min_{\overline{\Omega}} \phi'(u)} \leqslant c_3 e^{\gamma c_{12}^2} \text{ and we are done. Otherwise, we assume}$ that the maximum of Du is attained at an interior point p of Ω . Set

 $w(q) := \frac{u_x^2(q) + u_t^2(q)}{2}, q \in \Omega$. Thus, it suffices to infer the desired a priori estimates for the quantity $w(p) = \frac{u_x^2(p) + u_t^2(p)}{2}$. Of course, we have

(33)
$$w_x(p) = w_t(p) = 0$$

and $u_x^2 u_{xx} = u_t^2 u_{tt}$ at p

We remark that for the deduction of the last equation we used that we are working only with two independent variables x, t. This will allow us to write the second derivatives appearing in (27), in term of the first derivatives (at p). This is a crucial step to achieve the desired global estimates making use of the present method.

Hereafter, we write ϕ, ϕ', ϕ'' . Next plug the expressions (31) and its derivatives with respect to x and t, respectively, in (27). Then rewrite the ϵ -horizontal minimal equation in the form:

(34)
$$(\phi^{2} + 2\phi'^{2}w + \epsilon)u_{xx} + (1 + 2\phi'^{2}w)u_{tt} - \phi'^{2}(u_{x}w_{x} + u_{t}w_{t})$$
$$+ \frac{\phi^{2}\phi''}{\phi'}2w - \frac{\phi''(\phi^{2} - 1)}{\phi'}u_{t}^{2} + \frac{\phi}{\phi'}(1 + 2w\phi'^{2}) - \frac{\phi\phi'^{2}}{\phi'}u_{t}^{2} + \frac{\phi''}{\phi'}\epsilon u_{x}^{2} = 0$$

Now note that by invoking elliptic regularity $g \in C^3(\Omega)$. In fact, g is analytic in Ω by Morrey's regularity theorem [31]. Set $a_{11} := \phi^2 + \phi'^2 u_t^2 + \epsilon$, $a_{22} = 1 + \phi'^2 u_x^2$ and $a_{12} = a_{21} = -\phi'^2 u_x u_t$. The following identity will be useful and its verification is easy to check.

$$(35) \quad (\phi^{2} + 2\phi'^{2}w + \epsilon)(u_{x}u_{xxx} + u_{t}u_{txx}) + (1 + 2\phi'^{2}w)(u_{x}u_{xtt} + u_{t}u_{ttt}) - \phi'^{2}(u_{x}^{2}w_{xx} + u_{t}^{2}w_{tt} + 2u_{x}u_{t}w_{xt}) = (\phi^{2} + \phi'^{2}u_{x}^{2} + \phi'^{2}u_{t}^{2} + \epsilon)(u_{x}u_{xxx} + u_{t}u_{txx}) + (1 + \phi'^{2}u_{x}^{2} + \phi'^{2}u_{t}^{2})(u_{x}u_{xtt} + +u_{t}u_{ttt}) - \phi'^{2}(u_{x}^{2}w_{xx} + u_{t}^{2}w_{tt} + 2u_{x}u_{t}w_{xt}) = = (\phi^{2} + \phi'^{2}u_{x}^{2} + \phi'^{2}u_{t}^{2} + \epsilon)(w_{xx} - u_{xx}^{2} - u_{xt}^{2}) + (1 + \phi'^{2}u_{x}^{2} + \phi'^{2}u_{t}^{2})(w_{tt} - u_{tt}^{2} - u_{xt}^{2}) - \phi'^{2}(u_{x}^{2}w_{xx} + u_{t}^{2}w_{tt} + 2u_{x}u_{t}w_{xt}) = a_{11}w_{xx} + a_{22}w_{tt} + 2a_{12}w_{xt} - (\phi^{2} + \phi'^{2}u_{t}^{2} + \epsilon)(u_{xx}^{2} + u_{xt}^{2}) - (\phi'^{2}u_{x}^{2})(u_{xx}^{2} + u_{xt}^{2}) - (1 + \phi'^{2}u_{x}^{2})(u_{xt}^{2} + u_{tt}^{2}) - \phi'^{2}u_{t}^{2}(u_{xt}^{2} + u_{tt}^{2}) = a_{11}w_{xx} + a_{22}w_{tt} + 2a_{12}w_{xt} - (\phi^{2} + \epsilon + 2\phi'^{2}w)(u_{xx}^{2} + u_{xt}^{2}) - (1 + 2\phi'^{2}w)(u_{tt}^{2} + u_{xt}^{2})$$

The conditions a_{ij} is positive and $D^2w(p)$ is negative read

(36)
$$(\phi^2 + 2\phi'^2 w + \epsilon)(u_x u_{xxx} + u_t u_{txx}) + (1 + 2\phi'^2 w)(u_x u_{xtt} + u_t u_{ttt}) - \phi'^2(u_x^2 w_{xx} + u_t^2 w_{tt} + 2u_x u_t w_{xt}) \leq 0 \quad \text{at} \quad p.$$

Now differentiate the P.D.E (34) with respect to x (respectively differentiate with respect to t) and multiply the resulting equation by u_x (respectively multiply by u_t). Adding the two equations thus obtained and using (33) and (36), we infer

(37)

$$8\phi'\phi''w^{2}(u_{xx}+u_{tt})+4\phi\phi'w\,u_{xx}-2\phi'^{2}u_{t}^{2}w+6\phi''\phi u_{x}^{2}w+2w\left(\frac{\phi'^{2}-\phi''\phi}{\phi'^{2}}\right)$$

$$+4w^{2}\left[-\phi^{2}\left(\frac{-\phi'''\phi'+\phi''^{2}}{\phi'^{2}}\right)+\phi'^{2}\right]+2u_{t}^{2}w\left[(\phi^{2}-1)\left(\frac{-\phi'''\phi'+\phi''^{2}}{\phi'^{2}}\right)\right]$$

$$-\left(\frac{-\phi'''\phi'+\phi''^{2}}{\phi'^{2}}\right)\epsilon\,2wu_{x}^{2} \ge 0 \quad \text{at} \quad p.$$

Hence, according to the foregoing we find

(38)
$$(8\phi'\phi''w^2)(u_{xx} + u_{tt}) + (4\phi\phi'w)u_{xx}$$

 $+ 4w^2 \left[-\min\{\phi^2(u), 1\}\left(\frac{-\phi'''\phi' + \phi''^2}{\phi'^2}\right) + \phi'^2\right]$
 $+ (6\phi''\phi u_x^2w) + 2w\left(\frac{\phi'^2 - \phi''\phi}{\phi'^2}\right) \ge 0 \text{ at } p.$

since $\frac{-\phi'''\phi'+\phi''^2}{\phi'^2} = 2\gamma > 0$, noticing that the last term of inequality (37) is negative.

We must now estimate the quantities $u_{xx}(p)$ and $(u_{xx} + u_{tt})(p)$ in terms of ϕ, ϕ', ϕ'' and w at p. Owing to (33) performing some computations we find that (34) becomes

(39)
$$u_{xx} \phi' \left[(\phi^2 + \epsilon + 2\phi'^2 w) u_t^2 + (1 + 2\phi'^2 w) u_x^2 \right]$$

= $-u_t^2 \phi'' \phi^2 u_x^2 - \phi'' u_t^4 - \phi u_t^2 - \phi'^2 \phi u_t^2 u_x^2 - \phi'' \epsilon u_x^2 u_t^2$ at p

and

(40)
$$(u_{xx} + u_{tt}) \phi' \left[(\phi^2 + \epsilon + 2\phi'^2 w) u_t^2 + (1 + 2\phi'^2 w) u_x^2 \right]$$

= $-2\phi'' \phi^2 w u_x^2 - \phi'' u_t^2 2w - 2\phi w - 2\phi \phi'^2 w u_x^2 - 2\phi'' \epsilon w u_x^2$ at p .

We now perform some calculations to discover

(41)
$$(8\phi'\phi''w^2)(p)(u_{xx}+u_{tt})(p) \leqslant -\frac{4\phi\phi''w}{\phi'^2}(p) - (4\phi\phi''u_x^2w)(p)$$

and

$$(4\phi\phi'w)(p) u_{xx}(p) \leqslant -\frac{4\phi^3\phi''w}{\phi'^2}(p) - \frac{8\phi\phi''w}{\phi'^2}(p)$$

since $\phi' > 0$ and $\phi'' < 0$.

Inserting (41) into (38) we then obtain

(42)
$$4w^{2}(p) \left[-\min\{\phi^{2}(u(p)), 1\} \left(\frac{-\phi'''\phi' + \phi''^{2}}{\phi'^{2}} \right)(p) + \phi'^{2}(p) \right] + 2w(p) \left[\left(\frac{\phi'^{2} - \phi''\phi}{\phi'^{2}} \right)(p) - \frac{6\phi\phi''}{\phi'^{2}}(p) - \frac{2\phi^{3}\phi''}{\phi'^{2}}(p) \right] \ge 0$$

Employing (30) into (42) we at last infer

(43)
$$4w^{2}(p) \left[-2\gamma \min\{c_{1}^{2},1\}+1\right] + 2w(p) \left[1+\left((\gamma+2\gamma c_{12})c_{2}+(2\gamma+4\gamma c_{12})(3c_{2}+c_{2}^{3})\right)e^{2\gamma c_{12}+2\gamma c_{12}^{2}}\right] \ge 0$$

since $c_{1} > 0$.

But then,

(44)
$$w^{2}(p) \left[-2\gamma c_{1}^{2}+1\right]$$

 $+2w(p) \left[1+\left((\gamma+2\gamma c_{12})c_{2}+(2\gamma+4\gamma c_{12})(3c_{2}+c_{2}^{3})\right)e^{2\gamma c_{12}+2\gamma c_{12}^{2}}\right] \ge 0$

since we assume that $c_1 \leq 1$.

Finally, we recall that we have chosen γ , so that $2\gamma > \frac{1}{c_1^2}$. Henceforth, in the light of (44) we derive the a priori bounds for |Du(p)|, if $c_1 \leq 1$.

Now assume that $c_1 > 1$. Choose $\lambda < 1$ such that $\lambda c_1 < 1$. Write by $\overline{y} = g(\overline{x}, t)$, the solution of (27) in Ω , taking coordinates $(\overline{x}, \overline{y}, t) \in \mathbb{H}^2 \times \mathbb{R}$.

By employing the translations $(\overline{x}, \overline{y}, t) \to (\lambda \overline{x}, \lambda \overline{y}, t) := (x, y, t)$, we obtain another function $y = g_{\lambda}(x, t), (x, t) \in T_{\lambda}(\Omega)$ (Remark 2.2 (3)), satisfying the equation (27) in $T_{\lambda}(\Omega)$ for $\epsilon \lambda^2$ in the place of ϵ . In fact, noticing that $g_{\lambda}(x, t) = \lambda g(\overline{x}, t) = \lambda g(\frac{x}{\lambda}, t)$. It suffices to compute the relations between the first and second derivatives of $g(\overline{x}, t)$ and $g_{\lambda}(x, t)$.

We conclude therefore that g_{λ} satisfies

(45)

$$(g_{\lambda})_{xx}((g_{\lambda})^{2} + (g_{\lambda})_{t}^{2} + \epsilon\lambda^{2}) + (g_{\lambda})_{tt}(1 + (g_{\lambda})_{x}^{2}) - 2(g_{\lambda})_{x}(g_{\lambda})_{t}(g_{\lambda})_{xt} + (g_{\lambda})(1 + (g_{\lambda})_{x}^{2}) = 0$$

Now the same relations between the first derivatives of $g(\overline{x},t)$ and $g_{\lambda}(x,t)$, ensures that $\lambda^2(g_{\overline{x}}^2(\frac{x}{\lambda},t)+g_t^2(\frac{x}{\lambda},t)) \leq (g_{\lambda})_x^2(x,t)+(g_{\lambda})_t^2(x,t)$ and $(g_{\lambda})_x^2(x,t)+(g_{\lambda})_t^2(x,t) \leq g_{\overline{x}}^2(\frac{x}{\lambda},t)+g_t^2(\frac{x}{\lambda},t)$.

Thus

 $\max_{\partial T_{\lambda}(\Omega)} |Dg_{\lambda}| \leq \max_{\partial \Omega} |Dg| \text{ and } \max_{\overline{\Omega}} |Dg| \leq \frac{\max_{\overline{T_{\lambda}(\Omega)}} |Dg_{\lambda}|}{\lambda}.$

In view of $\lambda c_1 \leq g_\lambda \leq \lambda c_2$, in $T_\lambda(\Omega)$, we can commence again (since $\lambda c_1 < 1$), to mimic each step of the the above procedure to obtain the desired a priori bounds, if $c_1 > 1$.

Henceforth, the a priori global gradient estimates for |Dg| is achieved, as desired. This completes the proof of the theorem.

In the case of a convex domain, we state the following global C^1 estimates. First, if n = 2, equation (22) becomes (recall that $0 \leq \ell(s) \leq 1, s \in [0, 1], \ell(1) = 1, \ell(s) = 0, s \in [0, 1/2]$).

(46)
$$g_{xx}(g^2 + g_t^2 + \epsilon) + g_{tt}(1 + g_x^2) - 2g_xg_tg_{xt} + \ell(s)g(1 + g_x^2) = 0,$$

THEOREM 5.2 (Uniform global gradient estimates II).

Let $\Omega \subset \partial_{\infty} \mathbb{H}^2 \times \mathbb{R}$ be a $C^{2,\alpha}$ bounded convex domain for some $0 < \alpha < 1$. Let $\Gamma = \partial \Omega$ and let $f \in C^{2,\alpha}(\Gamma)$ be a positive function. Let $\epsilon \in [0,1]$. Given $s \in [1/2,1]$, let $g_s \in C^{2,\alpha}(\overline{\Omega})$ be a positive solution of equation (46) in Ω , such that $g_s = (2s-1)f + 2(1-s)\min_{\Gamma} f$ on Γ . Assume further that $R(\Omega, f) < \min_{\Gamma} f + \min_{\Gamma} f \sqrt{\frac{\pi}{2}}$, where $R(\Omega, f)$ is given by Definition 3.1 and Lemma 3.1.

Then, the following estimate holds.

(47)
$$\max_{\overline{\Omega}} |Dg_s| \leqslant C$$
where $C = C(\min_{\Gamma} f, \max_{\Gamma} f, \max_{\Omega} x - \min_{\Omega} x, K)$

where K is given by Lemma 4.1.

Proof. We summarize the proof as follows. In view of Theorem 4.2, it is routine to check that the proof of the global gradient (Theorem 5.1) is valid for equation (46), taking $c_2 = R(\Omega, f)$ and $c_1 = \min_{\Gamma} f$, noticing that $0 \leq \ell(s) \leq 1$. Indeed, we obtain the same a priori bounds as given by equations (43) and (44).

6. An existence result

Our main goal now is to prove the existence of a solution of a Dirichlet problem for the horizontal minimal equation. However, the non strictly ellipticity of the equation imposes a new insight.

Because of that, we need to consider the family of ϵ -horizontal minimal equation (27) ($0 \leq \epsilon \leq 1$) in order to solve the Dirichlet Problem for $\epsilon > 0$; then, by a compactness argument, we solve the original horizontal minimal equation.

Consider the following Dirichlet problem in two independent variables.

(48)
$$\mathcal{M}_{\mathcal{H}\epsilon}(g) = 0 \quad \text{in} \quad \Omega$$
$$g = f \quad \text{on} \quad \Gamma, \quad g \in C^{2,\,\alpha}(\bar{\Omega})$$

Let $R(\Omega, f)$ be the quantity defined in Definition 3.1. We have the following existence result.

THEOREM 6.1. Let $\Omega \subset \partial_{\infty} \mathbb{H}^2 \times \mathbb{R}$ be a $C^{2,\alpha}$ bounded convex domain for some $0 < \alpha < 1$. Let $\Gamma = \partial \Omega$ and let $f \in C^{2,\alpha}(\Gamma)$ be a positive function.

Assume further that $R(\Omega, f) < \min_{\Gamma} f + \min_{\Gamma} f \sqrt{\frac{\pi}{2}}$.

Then, for any $\epsilon \in [0, 1]$, the Dirichlet problem (48) admits a positive solution g.

Particularly, f admits an extension $g \in C^{2,\alpha}(\overline{\Omega})$ satisfying the horizontal minimal equation (4) on Ω . Furthermore, this solution is obtained as the limit in the C²-topology of a sequence $g_{\epsilon_n}, 0 < \epsilon_n < 1$ satisfying (48), as $\epsilon_n \to 0$.

We observe that, as in Theorem 5.1, the last assumption in Theorem 6.1 is invariant by hyperbolic translations.

Proof. Suppose first $0 < \epsilon \leq 1$. We first intend to show the existence of a solution of the Dirichlet problem (48), for $0 < \epsilon \leq 1$.

Let $\ell(s) = 2s - 1$, $s \in [1/2, 1]$ and $\ell(s) = 0$, $s \in [0, 1/2]$. Let $h(s, p) = (2s - 1)f(p) + 2(1 - s) \min_{\Gamma} f$, $s \in [1/2, 1]$ and $h(s, p) = 2s \min_{\Gamma} f$, $s \in [0, 1/2]$.

Given $s \in [0, 1]$ and given $g \in C^{1, \beta}(\overline{\Omega})$, consider the following family of linear Dirichlet problems:

$$\begin{split} u_{xx}(g^2 + g_t^2 + \epsilon) + u_{tt}(1 + g_x^2) - 2g_x g_t u_{xt} + \ell(s) \max\{g, 0\}(1 + g_x^2) &= 0\\ &\text{in} \quad \Omega,\\ u(p) &= h(s, p), \quad p \in \Gamma. \end{split}$$

Define an operator $T : C^{1,\beta}(\overline{\Omega}) \times [0,1] \to C^{2,\alpha\beta}(\overline{\Omega}), u = T(g, s)$ by the unique solution of the above problem for given (g, s). Existence and uniqueness of u are ensured by the extension lemma [17, Lemma 6.38] and by the classical theorem for linear strictly elliptic operator [17, Theorem 6.14]. Note that T(g,0) = 0 and $T(g,s) = 2s \min_{\Gamma} f$, if $s \in [0, 1/2]$.

To ensure the existence of a solution of our Dirichlet problem in the space $C^{2,\alpha}(\bar{\Omega})$ it is suffices to check that $T(\cdot, 1)$ has a fixed point. The equation g = T(g, s) reads

(49)

$$g_{xx}(g^2 + g_t^2 + \epsilon) + g_{tt}(1 + g_x^2) - 2g_x g_t g_{xt} + \ell(s) \max\{g, 0\}(1 + g_x^2) = 0$$
in Ω ,
 $g = h(s, \cdot)$ on Γ .

Then, we can apply maximum principle, comparing with Euclidean planes y = cst, to ensure that a solution of the equation (49) is nonnegative, so it satisfies

(50)
$$g_{xx}(g^2 + g_t^2 + \epsilon) + g_{tt}(1 + g_x^2) - 2g_x g_t g_{xt} + \ell(s)g(1 + g_x^2) = 0$$

(50) in Ω ,
 $g = h(s, \cdot)$ on Γ .

with $g \ge 0$ on Ω .

Noticing that by applying the maximum principle we have if $s \neq 0$, g > 0 on Ω and if s = 0, $g \equiv 0$, *i.e.* g satisfies equation (46), for $0 < \epsilon \leq 1$. Of course, the definition of $\ell(s)$ ensures that the solution g for $0 \leq s \leq 1/2$, is constant equal to $h(s, p) = 2s \min_{\Gamma} f$. Observe that

the solution g for $s \ge 1/2$, satisfies the inequality $g \ge \min_{\Gamma} f$ on Ω . Now in virtue of our a priori global C^1 estimates (Theorem 5.2) and the global Hölder estimates of Ladyzhenskaya and Ural'tseva [17], we have a priori global Hölder estimates for the first derivatives. That is, there exists a constant C such that $[Dg]_{\Omega,\beta} \le C$, for all g satisfying (50). Hence, by employing the Leray-Schauder theorem [2], [17], we obtain the desired existence of a positive solution g_{ϵ} of the Dirichlet problem (48), if $0 < \epsilon \le 1$. Recall that the uniform a priori horizontal length estimates, given by Lemma 3.1, forces the uniform lower bound $g_{\epsilon} > \min_{\Gamma} f$ on Ω , independently of ϵ .

Now let ϵ_n be a sequence such that $\epsilon_n \to 0$, if $n \to \infty$ $(0 < \epsilon_n \leq 1)$ and let g_{ϵ_n} be a positive $C^{2,\alpha}(\overline{\Omega})$ solution of (48). Our a priori global C^1 estimates combined with the a priori Schauder global estimates, allow us to apply the Arzelà-Ascoli's theorem to obtain a subsequence $\{g_{\epsilon_{n_j}}\}$ that converges to a $C^2(\overline{\Omega})$ nonnegative function g satisfying (27). Clearly, $g \ge \min_{\Gamma} f$ on Ω . Henceforth we have a solution of the Dirichlet problem (48), for $\epsilon = 0$. This accomplishes the proof of the theorem. \Box

REMARK 6.1. The geometry of the ambient space $\mathbb{H}^2 \times \mathbb{R}$ has some very intriguing geometric phenomenon, as we have commented in the introduction: Any catenoid (minimal surface of revolution) has vertical height less than π and the supremum of the family is π . E. Toubiana and the author, using the family of catenoids as suitable barriers, proved an asymptotic principle [45, Theorem 2.1] that have many consequences. In particular, it follows that there is no horizontal minimal graph given by a function $g \in C^2(\Omega) \cap C^0(\overline{\Omega})$ on a bounded strictly convex domain Ω , taking zero boundary data on $\partial\Omega$. We believe that the fact that the strict convexity of the Jordan domain forbids the existence of the Dirichlet problem for (4), with zero boundary data, is a very surprising phenomenon.

Some other results in the minimal surfaces theory, make use of the behavior of the catenoid family [34], [21]. The family of *n*-dimensional catenoids in $\mathbb{H}^n \times \mathbb{R}$ has an analogous interesting geometric behavior [6].

Finally, we remark that in [35] it is proved an Asymptotic Theorem setting in higher dimensions the non-existence result mentioned above: There is no horizontal minimal graph in $\mathbb{H}^n \times \mathbb{R}$ over a bounded strictly convex domain, given by a positive function g continuous up to the boundary, taking zero boundary value data.

On the other hand, the following existence result is somehow a counterpart of the above remark and is a immediate consequence of Theorem 6.1. Recall that $h(\Gamma) = \max_{\Gamma} x|_{\Gamma} - \min_{\Gamma} x|_{\Gamma}$ (Definition 3.1).

COROLLARY 6.1. Let $\Omega \subset \partial_{\infty} \mathbb{H}^2 \times \mathbb{R}$ be a $C^{2,\alpha}$ bounded convex domain for some $0 < \alpha < 1$. Let $\Gamma = \partial \Omega$ and let $f \in C^{2,\alpha}(\Gamma)$ be a positive function.

Let c_0 be a constant satisfying the inequality $c_0 \ge \operatorname{osc}_{\Gamma}(f) + \frac{h(\Gamma)}{2}$. Then, for any $\epsilon \in [0, 1]$, there exists a solution of the ϵ -horizontal minimal equation taking the boundary data $f + c_0$ on Γ .

References

- A. J. Aiolfi and C. V. Mathias, Existence and uniqueness of CMC parabolic graphs in H³ with boundary data satisfying the bounded slope condition, Diff. Geom. Appl. (6) 27,755-765 (2009).
- [2] H. Amann, "Lectures on some fixed point theorems", Monografias de Matemática, IMPA (1974).
- [3] J. Bakel'man, Hypersurfaces with given mean curvature and quasilinear elliptic equations with strong singularities, Mat. Sbornik (4) (75) 117, 561-595 (1968).
- [4] A. M. Barbosa. Compact minimal vertical graphs with non-connected boundary in Hⁿ × R,, Illin. J. Math. (3) 58, 593-617 (2014).
- [5] L. Barbosa and R. Sa Earp, Prescribed mean curvature hypersurfaces in Hⁿ⁺¹, with convex planar boundary, II, Sém. Th. Spect. et Geom. de Grenoble, 16, 43-79 (1998).
- [6] P. Bérard and R. Sa Earp, Minimal hypersurfaces in Hⁿ × R, total curvature and index, Bollettino dell'Unione Matematica Italiana. First online: 15 January 2016. Doi: 10.1007/s40574-015-0050-0.
- [7] P. Bérard, R. Sa Earp, Examples of H-hypersurfaces in Hⁿ × ℝ and geometric applications, Mat. Contem. 34, 19-51 (2008).
- [8] E. Bombieri, E. De Giorgi and M. Miranda, Una maggiorazione a priori relativa alle ipersuperfici minimali non parametriche, Arch. Rational Mech. Anal. 32, 255-267 (1969).
- [9] L. Caffarelli, L. Nirenberg and J. Spruck, Non-linear second order elliptic equations V. The Dirichlet problem for Weingarten hypersurfaces, Comm. on Pure and Applied Math 61, 47-70 (1988).
- [10] G. Citti and C. Senni, Constant mean curvature graphs on exterior domains of the hyperbolic plane, Math. Zeit. (1-2) 272, 531-550 (2012).
- [11] M. Dajczer and J. Ripoll, An extension of a theorem of Serrin to graphs in warped products, J. Geometric. Analysis (2) 15, 193-205 (2005).
- [12] M. F. Elbert, B. Nelli and R. Sa Earp, Existence of vertical ends of mean curvature 1/2 in H²×ℝ, Trans. Amer. Math. Soc. (3) 364, 1179-1191 (2012).

- [13] M. F. Elbert and R. Sa Earp, All solutions of the CMC-equation in Hⁿ × ℝ invariant by parabolic screw motion, Annali di Matematica Pura ed Applicata (1) 193, 103-114 (2014).
- [14] M. F. Elbert and R. Sa Earp, Constructions of H_r hypersurfaces, barriers and Alexandrov theorem in $\mathbb{H}^n \times \mathbb{R}$, Annali di Matematica Pura ed Applicata, **194**, 1809-1834 (2015).
- [15] R. Finn, On equations of minimal surface type, Annals of Math. 60, 397-416 (1954).
- [16] J. A. Gálvez and H. Rosenberg, Minimal surfaces and harmonic diffeomorphisms from the complex plane onto certain Hadamard surfaces, Amer. J. Math. (5) 132, 1249-1273 (2010).
- [17] D. Gilbarg and N.S. Trudinger, "Elliptic partial differential equations of second" order, Springer-Verlag (2001).
- [18] A. Granas and J. Dugundji, "Fixed point theory", Springer-Verlag (2010).
- [19] E. Guio and R. Sa Earp, Existence and non-existence for a mean curvature equation in hyperbolic space, Comm. Pure Appl. An. (3) 4, 549-568 (2005). Errata: Comm. Pure Appl. An. (2) 7, 465-465 (2008).
- [20] L. Hauswirth, H. Rosenberg & J. Spruck, On Complete Mean Curvature $H = \frac{1}{2}$ surfaces in $\mathbb{H}^2 \times \mathbb{R}$, Comm. Anal. Geom. (5) **16**, 989-1005 (2009).
- [21] L. Hauswirth, B. Nelli, R, Sa Earp and E. Toubiana, A Schoen theorem for minimal surfaces in H² × ℝ. Advances in Mathematics, 274, 199-240 (2015). doi: 10.1016/j.aim.2014.12.030.
- [22] H. Jenkins and J. Serrin, Variational problems of minimal surface type I, Arch. Rat. Mech. Ana. 12, 185-212 (1963).
- [23] H. Jenkins and J. Serrin, The Dirichlet problem for the minimal surface equation in higher dimensions, J. Reine Angew. Math. 229, 170-187 (1968).
- [24] O. A. Ladyzhenskaya and N. N. Ural'tseva, "Linear and quasilinear elliptic equations" (English translation), New York, Academic Press 1968. 2nd Russian ed. (1973).
- [25] H. F. Lima and U. L. Parente, A Bernstein type theorem in $\mathbb{R} \times \mathbb{H}^n$, Bull. Braz. Math. Soc. New Series (1) 43, 17-26 (2012).
- [26] R. López and S. Montiel. Existence of constant mean curvature graphs in hyperbolic space, Calc. Var. 8, 177-190 (1999).
- [27] F. Martin, R. Mazzeo, M. Rodríguez. Minimal surfaces with positive genus and finite total curvature in H²×ℝ. Geometry & Topology 18, 141-177 (2014).
- [28] L. Mazet, M. Rodriguez and H. Rosenberg, The Dirichlet problem for the minimal surface equation - with possible infinite boundary data - over domains in a Riemannian surface, Proc. London Math. Soc. (6) 102, 985-1023 (2011).
- [29] L. Mazet, M. Rodriguez and H. Rosenberg, *Periodic constant mean curvature surfaces in* H² × ℝ, Asian J. Math (5) **18**, 829-858 (2014).
- [30] C. B. Morrey, The problem of Plateau on a Riemannian manifold, Ann. Math. No. 2, 49, 807-851 (1948).

- [31] C. B. Morrey, "Multiple integrals in the calculus of variations", Reprint of the 1966 Edition, Springer (2008).
- [32] B. Nelli and J. Spruck, "On the existence and uniqueness of constant mean curvature hypersurfaces in hyperbolic space", In: Geometric analysis and calculus of variations, Intern. Press, 253-266 (1996).
- [33] B. Nelli and R. Sa Earp, Some properties of hypersurfaces of prescribed mean curvature in ℍⁿ⁺¹, Bull. Sci. Math. **120**, 537-553 (1996).
- [34] B. Nelli, R. Sa Earp, W. Santos, E. Toubiana, Uniqueness of H-surfaces in $\mathbb{H}^2 \times \mathbb{R}$, $|H| \leq 1/2$, with boundary one or two parallel horizontal circles, Ann. Global Anal. Geom. (4) **33**, 307-321 (2008).
- [35] B. Nelli, R. Sa Earp and E. Toubiana, Maximum Principle and Symmetry for Minimal Hypersurfaces in Hⁿ × ℝ, Annali della Scuola Normale Superiore di Pisa, Classe di Scienze (2015). doi: 10.2422/2036 – 2145.201211_004.
- [36] B. Nelli and H. Rosenberg, *Minimal surfaces in* H² × ℝ, Bull. Braz. Math. Soc., New Series (2) **33**, 263-292 (2002); Errata: Bull. Braz. Math. Soc., New Series (4) **38**, 1-4 (2007).
- [37] B. Nelli, R. Sa Earp and E. Toubiana, Minimal Graphs in Nil₃: existence and non-existence results. arXiv: 1508.01724 [math.DG] (2015).
- [38] L. Nirenberg, "Topics in nonlinear functional analysis", Courant Lecture Notes in Mathematics. American Mathematical Society (2001).
- [39] M. H. Protter and H. F. Weinberger, "Maximum principles in differential equations", Prentice-Hall (1967).
- [40] P. Pucci and J. B. Serrin, "The maximum principle" (Progress in nonlinear differential equations and their applications), Birkhäuser (2007).
- [41] H. Rosenberg, R. Souam, E. Toubiana, General Curvature Estimates for Stable H-surfaces in 3-manifolds and Applications, J. Diff. Geom. 84, 623-648 (2010).
- [42] R. Sa Earp, Parabolic and Hyperbolic Screw motion in H²×ℝ, Journ. Austra. Math. Soc 85, 113-143 (2008).
- [43] R. Sa Earp, Uniqueness of minimal surfaces whose boundary is an horizontal graph and the Bernstein problems in H² × ℝ, Math. Zeit. (1) 273, 211-217 (2013).
- [44] R. Sa Earp and E. Toubiana, Existence and uniqueness of minimal graphs in hyperbolic space, Asian J. Math. 4, 669-694 (2000).
- [45] R. Sa Earp and E. Toubiana, An asymptotic theorem for minimal surfaces and existence results for minimal graphs in H² × ℝ, Math. Annalen (2) 342, 309-331 (2008).
- [46] R. Sa Earp and E. Toubiana, "Introduction à la géométrie hyperbolique et aux surfaces de Riemann", Cassini (2009).
- [47] R. Sa Earp and E. Toubiana, *Minimal graphs in* $\mathbb{H}^n \times \mathbb{R}$ and \mathbb{R}^{n+1} , Ann. de l'Inst. Fourier (7) **60**, 2373-2402 (2010).

- [48] L. Simon, Equations of Mean Curvature Type in 2 Independent Variables, Pac. Journ. of Math. (1) 69, 245-268 (1977).
- [49] J. Spruck, Interior Gradient Estimates and Existence Theorems for Constant Mean Curvature Graphs in Mⁿ × ℝ, Pure Appl. Math. Q. (3) part 2, 785-800 (2007).

Departamento de Matemática, Pontifícia Universidade Católica do Rio de Janeiro, Rio de Janeiro, 22453-900 RJ, Brazil

E-mail address: rsaearp@gmail.com