

Dynamics of the symmetric eigenvalue problem with shift strategies

Ricardo S. Leite, Nicolau C. Saldanha and Carlos Tomei

August 23, 2011

Abstract

A common algorithm for the computation of eigenvalues of real symmetric tridiagonal matrices is the iteration of certain special maps F_σ called *shifted QR steps*. Such maps preserve spectrum and a natural common domain is \mathcal{T}_Λ , the manifold of real symmetric tridiagonal matrices conjugate to the diagonal matrix Λ . More precisely, a (generic) *shift* $s \in \mathbb{R}$ defines a map $F_s : \mathcal{T}_\Lambda \rightarrow \mathcal{T}_\Lambda$. A strategy $\sigma : \mathcal{T}_\Lambda \rightarrow \mathbb{R}$ specifies the shift to be applied at T so that $F_\sigma(T) = F_{\sigma(T)}(T)$. Good shift strategies should lead to fast *deflation*: some off-diagonal coordinate tends to zero, allowing for reducing of the problem to submatrices. For topological reasons, continuous shift strategies do not obtain fast deflation; many standard strategies are indeed discontinuous. Practical implementation only gives rise systematically to *bottom deflation*, convergence to zero of the lowest off-diagonal entry $b(T)$. For most shift strategies, convergence to zero of $b(T)$ is cubic, $|b(F_\sigma(T))| = \Theta(|b(T)|^k)$ for $k = 3$. The existence of arithmetic progressions in the spectrum of T sometimes implies instead quadratic convergence, $k = 2$. The complete integrability of the Toda lattice and the dynamics at non-smooth points are central to our discussion. The text does not assume knowledge of numerical linear algebra.

Keywords: Isospectral manifold, Deflation, Wilkinson's shift, *QR* algorithm.

MSC2010-class: 37N30, 37C05, 65F15.

1 Introduction

In this paper, we study some subtle dynamical aspects of a class of numerical algorithms for eigenvalues of real symmetric matrices. This includes the classic inverse iteration with different shift strategies, among which Rayleigh and Wilkinson shifts. We do not assume previous knowledge of numerical linear algebra.

Numerical analysts are familiar with *tridiagonalization*, the fact that given a real symmetric matrix S it is easy to obtain another isospectral matrix T which is tridiagonal: $(T)_{ij} = 0$ whenever $|i - j| > 1$. For matrices of order approximately between 20 and 1000, it pays to first tridiagonalize and then work in the vector space \mathcal{T} of real symmetric tridiagonal matrices. Let $\Lambda = \text{diag}(\lambda_1 < \lambda_2 < \dots < \lambda_n)$ be a diagonal matrix with simple spectrum: it turns out that the set $\mathcal{T}_\Lambda \subset \mathcal{T}$ of tridiagonal matrices isospectral with Λ is a connected compact smooth oriented manifold ([16], [8]). The algorithms under consideration are defined by iteration of some easily computable map $F : \mathcal{T}_\Lambda \rightarrow \mathcal{T}_\Lambda$: given $T \in \mathcal{T}_\Lambda$ we consider the sequence $(F^k(T_0))$. For relevant maps F , diagonal matrices in \mathcal{T}_Λ are fixed points of F .

Let $\mathcal{E} \subset O(n)$ be the group of real orthogonal diagonal matrices, so that for $E \in \mathcal{E}$, $(E)_{ii} = \pm 1$. For each $E \in \mathcal{E}$, the map $\eta : \mathcal{T}_\Lambda \rightarrow \mathcal{T}_\Lambda$, $\eta(T) = ETE$, is an involutive diffeomorphism of \mathcal{T}_Λ : its effect on $T \in \mathcal{T}_\Lambda$ is to change signs of some

subdiagonal entries $(T)_{i+1,i}$. Numerical analysts, familiar with this simple fact, often drop signs of subdiagonal entries. We shall not do likewise for we are often interested in smoothness issues. Again, relevant maps will be (\mathcal{E}) -equivariant, in the sense that $F \circ \eta = \eta \circ F$ for all η .

Consistently with the involutions above, signs of subdiagonal entries induce a cell decomposition of \mathcal{T}_Λ . The 0-cells are the $n!$ diagonal matrices and the top dimensional $(n-1)$ -cells turn out to be 2^{n-1} permutohedra (polytopes equivalent to the convex hull of the $n!$ points of \mathbb{R}^n obtained by permuting n fixed distinct real numbers). For $n=3$, the manifold \mathcal{T}_Λ is a bitorus which can be obtained by gluing four hexagons along six circles (see Figure 1).




Figure 1: The cell decomposition of \mathcal{T}_Λ for $\Lambda = \text{diag}(4, 5, 7)$

We consider that iteration of the map F has accomplished its job when one subdiagonal entry $(F^k(T))_{i+1,i}$ has absolute value smaller than some prescribed tolerance. In terms of the cell decomposition, we are done when we hit (a thin neighborhood of) a lower dimensional cell, or, in the numerical jargon, the sequence $(F^k(T))$ undergoes *deflation*. Notice that if $(T)_{i+1,i} = 0$ then the matrix T splits as $T = T^a \oplus T^b$ where the tridiagonal submatrices T^a and T^b have orders i and $n-i$, respectively. Pragmatically, if $(T)_{i+1,i} \approx 0$ then the spectrum of T is approximately the disjoint union of the spectra of T^a and T^b , which are easier to compute.

Ideally, deflation should happen approximately in the middle so that each subproblem has order approximately half of the original one. Unfortunately, it is not known how to implement easily computable iterations with this property. Usually the sequence $(F^k(T))$ undergoes *bottom deflation*:

$$\lim_{k \rightarrow +\infty} b(F^k(T)) = 0; \quad b(T) = (T)_{n,n-1}.$$

Geometrically, we approach one of the n deflation sets $\mathcal{D}_{\Lambda,0}^i \subset \mathcal{T}_\Lambda$ defined by $(T)_{n,n} = \lambda_i$, $b(T) = 0$.

Notice that removing the n -th row and column obtains a diffeomorphism:

$$\mathcal{D}_{\Lambda,0}^i \approx \mathcal{T}_{\Lambda_i}, \quad \Lambda_i = \text{diag}(\lambda_1, \dots, \lambda_{i-1}, \lambda_{i+1}, \dots, \lambda_n);$$

in particular, $\mathcal{D}_{\Lambda,0}^i$ is connected. In Figure 1, the submanifolds $\mathcal{D}_{\Lambda,0}^i$ are three of the six (removed) circles. It turns out (Proposition 4.2) that, for sufficiently small $\epsilon > 0$, the closed set $\mathcal{D}_{\Lambda,\epsilon}$ defined by $|b(T)| \leq \epsilon$ has n connected components $\mathcal{D}_{\Lambda,\epsilon}^i$ which are closed tubular neighborhoods of $\mathcal{D}_{\Lambda,0}^i$.

As algorithms for eigenvalue computation, continuous maps F are problematic.

Theorem 1 *Let $F : \mathcal{T}_\Lambda \rightarrow \mathcal{T}_\Lambda$ be a continuous \mathcal{E} -equivariant map such that every diagonal matrix in \mathcal{T}_Λ is a fixed point of F .*

(a) *The map F is surjective.*

(b) *If there exist disjoint compact sets $\mathcal{K}_i \supset \mathcal{D}_{\Lambda,0}^i$ with $F(\mathcal{K}_i) \subset \text{int}(\mathcal{K}_i)$ then there exists $T \in \mathcal{T}_\Lambda$ for which the sequence $(F^k(T))$ does not undergo bottom deflation.*

Item (a) already makes F unpromising as an algorithm: for any k , the iterate F^k is surjective and, given k , there exists T such that $F^k(T)$ is far from deflation. The additional hypothesis in item (b), which, as we shall see, holds for many algorithms, makes F even less desirable. The proof of this result uses methods very different from the rest of the paper and is left for the Appendix.

These phenomena lead numerical analysts to consider discontinuous maps F . Among the standard algorithms to compute eigenvalues of matrices in \mathcal{T} are QR steps with different shift strategies: Rayleigh and Wilkinson are familiar examples (excellent references are [17], [5], [14]). Recall that Rayleigh's strategy ρ is continuous and is known to have the unfortunate property (b) that there exists a matrix T for which $(F_\rho^k(T))$ does not undergo bottom deflation; Wilkinson, on the other hand, is discontinuous. In this paper, we consider a more general context: we define simple shift strategies, which include the examples above and more.

More precisely, given a matrix $T \in \mathcal{T}$ and $s \in \mathbb{R}$, write $T - sI = QR$, if possible, for an orthogonal matrix Q and an upper triangular matrix R with positive diagonal entries. A *shifted QR step* is $\Phi(T, s) = Q^*TQ$. As is well known, shifted QR steps preserve spectrum and shape. A function $\sigma : \mathcal{T}_\Lambda \rightarrow \mathbb{R}$ is (\mathcal{E} -)invariant if $\sigma(\eta(T)) = \sigma(ETE) = \sigma(T)$ for all $T \in \mathcal{T}_\Lambda$ and all $E \in \mathcal{E}$. A *simple shift strategy* is an invariant function $\sigma : \mathcal{T}_\Lambda \rightarrow \mathbb{R}$ satisfying the following condition: there exists $C_\sigma > 0$ such that for all $T \in \mathcal{T}_\Lambda$ there is an eigenvalue λ_i with $|\sigma(T) - \lambda_i| \leq C_\sigma |b(T)|$.

For technical reasons, we prefer the *signed step* $F_s(T) = \Phi_*(T, s) = Q_*^*TQ_*$, where now $T - sI = Q_*R_*$, the orthogonal matrix Q_* has positive determinant and only the first $n-1$ diagonal entries of the upper triangular matrix R_* are required to be positive. As we shall see, the signed step is smoothly defined on a larger domain, and convergence issues for both kinds of step iterations are essentially equivalent.

Simple shift strategies prescribe shifts: set $F_\sigma(T) = F_{\sigma(T)}(T)$. It turns out that F_σ is a well-defined (but usually discontinuous) equivariant map from \mathcal{T}_Λ (or some very large subset thereof) to \mathcal{T}_Λ .

An important question in practice is estimating the rate of deflation, i.e., the rate of convergence to zero of the sequence $b(F_\sigma^k(T))$. Numerical evidence indicates that deflation is often cubic, in the sense that there is a constant C such that $|b(F_\sigma^{k+1}(T))| \leq C|b(F_\sigma^k(T))|^3$ for large k .

Consider the *singular support* $\mathcal{S}_\sigma \subset \mathcal{T}_\Lambda$ of a shift strategy σ , the minimal closed subset of \mathcal{T}_Λ on whose complement σ is smooth. Away from the singular support \mathcal{S}_σ , squeezing is cubic.

Theorem 2 *For $\epsilon > 0$ small enough, each deflation neighborhood $\mathcal{D}_{\Lambda,\epsilon}^i$ is invariant under F_σ . There exists $C > 0$ such that, for all $T \in \mathcal{D}_{\Lambda,\epsilon}$, $|b(F_\sigma(T))| \leq C|b(T)|^2$. Also, given a compact set $\mathcal{K} \subset \mathcal{D}_{\Lambda,\epsilon}$ disjoint from $\mathcal{S}_\sigma \cap \mathcal{D}_{\Lambda,0}$, there exists $C_{\mathcal{K}} > 0$ such that, for all $T \in \mathcal{K}$, $|b(F_\sigma(T))| \leq C_{\mathcal{K}}|b(T)|^3$.*

Although the tubular neighborhoods $\mathcal{D}_{\Lambda,\epsilon}^i$ are invariant under F_σ , it is not true in general that $F_\sigma^k(T)$ belongs to $\mathcal{D}_{\Lambda,\epsilon}^i$ for sufficiently large k : this is true, however, for the important example of Wilkinson's shift.

For Rayleigh's shift, it is well known that convergence (when it happens) is always cubic; this is a corollary of Theorem 2. Cubic convergence does not hold in general for Wilkinson's strategy. In [9], for $\Lambda = \text{diag}(-1, 0, 1)$, we construct a Cantor-like set $\mathcal{X} \subset \mathcal{T}_\Lambda$ of unreduced initial conditions for which the rate of convergence is strictly quadratic. Sequences starting at \mathcal{X} converge to a reduced matrix which is not diagonal. A part of the set \mathcal{X} is shown (true to scale) on the left part of Figure 2; the Cantor-like aspect is invisible: the cross section of each of the four visible "curves" really consists of a tiny Cantor set, far smaller than the resolution of the picture. This is consistent with the fact that the Hausdorff dimension of \mathcal{X} is 1. The set \mathcal{X} is the intersection of thinner and thinner wedges; in the right half we show schematically three generations of such wedges. The central vertical line is a step-like discontinuity; the inverse image of the largest wedge \mathcal{X}_0 is $\mathcal{X}_1^{(+)} \cup \mathcal{X}_1^{(-)}$ and the inverse image of that is $\mathcal{X}_2^{(++)} \cup \mathcal{X}_2^{(+-)} \cup \mathcal{X}_2^{(-+)} \cup \mathcal{X}_2^{(--)}$.




Figure 2: The set \mathcal{X}

As numerical analysts know, shift strategies usually define sequences of matrices which, asymptotically, not only isolate an eigenvalue at the (n, n) position but also isolate, at a slower rate, a second eigenvalue at the $(n-1, n-1)$ position. This does not happen for the example above where $(F_\sigma^k(T))_{n,n}$ tends to the center of a three-term arithmetic progression of eigenvalues and $(F_\sigma^k(T))_{n-1, n-2}$ stays bounded away from zero.

A matrix $T \in \mathcal{T}$ with simple spectrum is *a.p. free* if it does not have three eigenvalues in arithmetic progression and *a.p. otherwise*; in particular, generic spectra are a.p. free. In this case, the situation is very nice: cubic convergence is essentially uniform on \mathcal{T}_Λ . This condition is reminiscent of the Sternberg's resonance hypothesis for normal forms ([15]).

Theorem 3 *Let Λ be an a.p. free matrix and σ a shift strategy for which diagonal matrices do not belong to \mathcal{S}_σ . Then there exist $\epsilon > 0$, $C > 0$ and $K > 0$ such that:*

- (a) *the deflation neighborhood $\mathcal{D}_{\Lambda, \epsilon}$ is invariant under F_σ ;*
- (b) *for any $T \in \mathcal{D}_{\Lambda, \epsilon}$, the sequence $(F_\sigma^k(T))$ converges to a diagonal matrix and the set of positive integers k for which $|b(F_\sigma^{k+1}(T))| > C|b(F_\sigma^k(T))|^3$ has at most K elements.*

Still, the finite set of points in which the cubic estimate does not hold may occur arbitrarily late along the sequence $(F_\sigma^k(T))$.

An a.p. matrix is *strong a.p.* if it contains three consecutive eigenvalues in arithmetic progression and *weak a.p.* otherwise. Under very mild additional hypothesis, $b(T)$ converges to zero at a cubic rate also for weak a.p. matrices. Let $\mathcal{C}_{\Lambda, 0} \subset \mathcal{T}_\Lambda$ be the set of matrices T for which $(T)_{n, n-1} = (T)_{n-1, n-2} = 0$.

Theorem 4 *Let Λ be a weak a.p. matrix and $\sigma : \mathcal{T}_\Lambda \rightarrow \mathbb{R}$ a shift strategy for which $\mathcal{C}_{\Lambda, 0}$ and \mathcal{S}_σ are disjoint. Then there exists $\epsilon > 0$ such that the deflation neighborhood $\mathcal{D}_{\Lambda, \epsilon}$ is invariant under F_σ and, for all unreduced $T \in \mathcal{D}_{\Lambda, \epsilon}$, the sequence $(b(F_\sigma^k(T)))$ converges to zero at a rate which is at least cubic. More precisely, for*

each unreduced $T \in \mathcal{D}_{\Lambda, \epsilon}$ there exist $C_T, K_T > 0$ such that, for all $k > K_T$, we have $|b(F_\sigma^{k+1}(T))| \leq C_T |b(F_\sigma^k(T))|^3$.

In particular, the convergence of Wilkinson's strategy is cubic for weak a.p. matrices. However, uniformity in the sense of Theorem 3 is not guaranteed and the constants C_T and K_T depend on T . As in the case of the spectrum $\{-1, 0, 1\}$, we conjecture that if Λ is strong a.p. then there exists $\mathcal{X} \subset \mathcal{T}_\Lambda$ of Hausdorff codimension 1 of initial conditions T for which the rate of convergence is strictly quadratic.

The celebrated integrability of the Toda lattice ([6], [13]) on unreduced tridiagonal matrices manifests itself in several ways along the paper: it provided ample inspiration but the paper strives to be self-contained. For starters, the steps F_s , $s \in \mathbb{R}$, commute in their natural domains (Proposition 2.6). Norming constants (as in [13]) provide angle variables for which steps F_s are translations. Unfortunately, these angle variables break down (as they must!) for reduced matrices $T \in \mathcal{T}_\Lambda$. Since $(F_\sigma^k(T))$ approaches reduced matrices we prefer to introduce other coordinate systems which extend smoothly to such points. *Bidiagonal coordinates*, defined in [8], consist of very explicit charts on the manifold \mathcal{T}_Λ . They are used in [8] to prove the cubic convergence of Rayleigh's shift and in the unpublished manuscript [10] to prove some of the results presented here for Wilkinson's shift. In Section 4, instead, we introduce *tubular coordinates* on the tubular neighborhoods $\mathcal{D}_{\Lambda, \epsilon}^i$: steps F_s within these sets are given by a very simple formula (Corollary 4.3).

The (signed) steps F_σ are smooth whenever the shift strategy is, i.e., for $T \in \mathcal{D}_{\Lambda, \epsilon}^i \setminus \mathcal{S}_\sigma$ ("unsigned steps" would not be smooth on limit points). At matrices $T_0 \in \mathcal{D}_{\Lambda, 0}^i$ on which F_σ is smooth, the map $T \mapsto b(F_\sigma(T))$ has zero gradient. The symmetry of the shift strategy yields a cubic Taylor expansion and therefore an estimate $|b(F_\sigma(T))| \leq C|b(T)|^3$, settling Theorem 2.

Height functions $H : \mathcal{D}_{\Lambda, \epsilon}^i \rightarrow \mathbb{R}$ (similar to Lyapunov functions) are used for further study of the sequence $(F_\sigma^k(T))$ in the a.p. free case. More precisely, for steps s near λ_i , $H_i(F_s(T)) > H_i(T)$ provided $T \in \mathcal{D}_{\Lambda, \epsilon}^i$ is not diagonal: this is another manifestation of the Toda dynamics. Theorem 3 then follows by a compactness argument bounding the number of iterations for which $F_\sigma^k(T)$ stays close to the singular support \mathcal{S}_σ .

For a.p. spectra the situation is subtler, as can be seen from the example in [9] and Figure 2. On the other hand, Theorem 4 tells us that the weak a.p. hypothesis together with an appropriate smoothness condition guarantee cubic convergence.

In Section 2 we list the basic properties of the signed shifted QR step on the manifold \mathcal{T}_Λ . Simple shift strategies are introduced in Section 3, and the standard examples are shown to satisfy the definition. We define the deflation set $\mathcal{D}_{\Lambda, 0}$ and neighborhood $\mathcal{D}_{\Lambda, \epsilon}$ in Section 4 and then set up tubular coordinates. The local theory of steps F_s near $\mathcal{D}_{\Lambda, 0}$ and the proof Theorem 2 are presented in Section 5. In Section 6 we construct the height functions H and then prove Theorem 3. The convergence properties for a.p. matrices in Theorem 4 are proved in Section 7. We present in Section 8 two counterexamples to natural but incorrect strengthenings of Theorems 3 and 4. Finally, the Appendix is dedicated to Theorem 1.

The authors are very grateful for the abundant contributions of several readers of this work and its previous versions. The authors acknowledge support from CNPq, CAPES, INCT-Mat and FAPERJ.

2 The manifold \mathcal{T}_Λ and shifted steps F_s

Let \mathcal{T} denote the real vector space of $n \times n$ real, symmetric, tridiagonal matrices endowed with the norm $\|T\|^2 = \text{tr}(T^2)$. For $T \in \mathcal{T}$, the *subdiagonal entries* of T are $(T)_{i+1, i}$ for $i = 1, \dots, n-1$. The lowest subdiagonal entry of T is $b(T) = (T)_{n, n-1}$.

As usual, let $SO(n)$ denote the set of orthogonal matrices with determinant equal to 1. Let Λ be a real diagonal matrix with simple eigenvalues $\lambda_1 < \dots < \lambda_n$. Define the *isospectral manifold*

$$\mathcal{T}_\Lambda = \{Q^* \Lambda Q, Q \in SO(n)\} \cap \mathcal{T},$$

the set of matrices in \mathcal{T} similar to Λ . The set $\mathcal{T}_\Lambda \subset \mathcal{T}$ is a real smooth manifold ([16]; [8] describes an explicit atlas of \mathcal{T}_Λ).

For a matrix M , the QR factorization is $M = QR$ for an orthogonal matrix Q and an upper triangular matrix R with positive diagonal. The $Q_\star R_\star$ factorization, instead, is $M = Q_\star R_\star$, for $Q_\star \in SO(n)$ and R_\star an upper triangular matrix with $(R_\star)_{i,i} > 0$, $i = 1, \dots, n-1$. A real $n \times n$ matrix M is *almost invertible* if its first $n-1$ columns are linearly independent: notice that almost invertible matrices are dense within $n \times n$ matrices and form an open set. The diagonal matrix E_{n-1} is such that $(E_{n-1})_{i,i}$ is 1 for $i < n$ and -1 for $i = n$.

Proposition 2.1 *An almost invertible real matrix M admits a unique $Q_\star R_\star$ factorization, with Q_\star and R_\star depending smoothly on M . If M is invertible, it admits unique (smooth) factorizations $M = QR = Q_\star R_\star$. If $\det M > 0$, the factorizations are equal, i.e., $Q = Q_\star$ and $R = R_\star$. If $\det M < 0$, $Q = Q_\star E_{n-1}$ and $R = E_{n-1} R_\star$. If $\det M = 0$, $(R_\star)_{n,n} = 0$.*

Proof: Let M be almost invertible. Applying Gram-Schmidt with positive normalizations on its first $n-1$ columns we obtain the first $n-1$ columns of both Q and R , as well as those of Q_\star and R_\star . The last column $v = Q_\star e_n$ of Q_\star is already well defined, by orthonormality and the fact that $\det Q_\star = 1$. Now, set $R_\star = M(Q_\star)^*$. The positivity of $R_{n,n}$ specifies whether the last column of Q is v or $-v$. Smoothness is clear by construction.

If M is invertible, $\det M = \det Q_\star \det R_\star$ implies that the last diagonal entry of R_\star has the same sign of $\det M$: the relations between the factorizations then follow. If M is not invertible, the relation among determinants implies $(R_\star)_{n,n} = 0$. ■

If all subdiagonal entries of T are nonzero, T is an *unreduced* matrix; otherwise, T is *reduced*. Notice that an unreduced tridiagonal matrix is almost invertible: indeed, the block formed by rows $2, \dots, n$ and columns $1, \dots, n-1$ is an upper triangular matrix with nonzero diagonal entries, and therefore, invertible.

We consider the *shifted QR step* and its *signed* counterpart,

$$\Phi(T, s) = Q^* T Q, \quad \Phi_\star(T, s) = Q_\star^* T Q_\star,$$

where $T - sI = QR$ and $T - sI = Q_\star R_\star$. Let $\text{Dom}(\Phi)$ be the set of pairs $(T, s) \in \mathcal{T} \times \mathbb{R}$ for which $T - sI$ is invertible: $\text{Dom}(\Phi)$ is open and dense in $\mathcal{T} \times \mathbb{R}$ and, from the Gram-Schmidt algorithm, Φ is smooth in $\text{Dom}(\Phi)$. Similarly, the above proof shows that Φ_\star is smooth in $\text{Dom}(\Phi_\star)$, with $(T, s) \in \text{Dom}(\Phi_\star)$ if $T - sI$ is almost invertible. Clearly, $\text{Dom}(\Phi)$ is strictly contained in $\text{Dom}(\Phi_\star)$.

Lemma 2.2 *For $(T, s) \in \text{Dom}(\Phi)$ (resp. $\text{Dom}(\Phi_\star)$), we have $\Phi(T, s) \in \mathcal{T}$ (resp. $\Phi_\star(T, s) \in \mathcal{T}$). The spectra of T , $\Phi(T, s)$ and $\Phi_\star(T, s)$ are equal. In the appropriate domains, for $T - sI = QR = Q_\star R_\star$ and $i = 1, 2, \dots, n-1$,*

$$(\Phi(T, s))_{i+1,i} = \frac{(R)_{i+1,i+1}}{(R)_{i,i}} (T)_{i+1,i}, \quad (\Phi_\star(T, s))_{i+1,i} = \frac{(R_\star)_{i+1,i+1}}{(R_\star)_{i,i}} (T)_{i+1,i}.$$

Thus, the top $n-2$ subdiagonal entries of T , $\Phi(T, s)$ and $\Phi_\star(T, s)$ have the same sign; also, $\text{sign}(T)_{n,n-1} = \text{sign}(\Phi(T, s))_{n,n-1}$.

Proof: We prove the statements for Φ_* ; the others are then easy.

For a pair $(T, s) \in \text{Dom}(\Phi) \subset \text{Dom}(\Phi_*)$, there are two expressions for $\Phi_*(T, s)$:

$$\Phi_*(T, s) = Q_*^* T Q_* = R_* T R_*^{-1}, \quad \text{where } T - sI = Q_* R_*.$$

From the first equality, $\Phi_*(T, s)$ is symmetric and from the second, $\Phi_*(T, s)$ is an upper Hessenberg matrix so that $\Phi_*(T, s) \in \mathcal{T}$ is similar to T . More generally, for $(T, s) \in \text{Dom}(\Phi_*)$ we still have

$$\Phi_*(T, s) = Q_*^* T Q_*, \quad \Phi_*(T, s) R_* = R_* T$$

and therefore $\Phi_*(T, s) \in \mathcal{T}$ is similar to T . Compute the $(i+1, i)$ entry of the second equation above to obtain $(\Phi_*(T, s))_{i+1, i} (R_*)_{i, i} = (R_*)_{i+1, i+1} (T)_{i+1, i}$, completing the proof. \blacksquare

The following result describes the behavior of Φ_* at points not in $\text{Dom}(\Phi)$, which will play an important role throughout the paper.

Lemma 2.3 *If $(T, s) \in \text{Dom}(\Phi_*) \setminus \text{Dom}(\Phi)$ then*

$$b(\Phi_*(T, s)) = (\Phi_*(T, s))_{n, n-1} = 0, \quad (\Phi_*(T, s))_{n, n} = s.$$

At a point $(T, s) \in \text{Dom}(\Phi_)$ with $b(T) = 0$ and $s = (T)_{n, n}$ we have $\text{grad}(b \circ \Phi_*) = 0$.*

Proof: Since $T - sI = Q_* R_* = R_*^* Q_*^*$ is not invertible then $(R_*)_{n, n} = 0$ and therefore $R_*^* e_n = 0$. Thus $v = (Q_*^*)^{-1} e_n = Q e_n$ satisfies $(T - sI)v = 0$. We then have $\Phi_*(T, s)e_n = Q^* T Q e_n = Q^* T v = Q^*(sv) = s e_n$, proving the first claim. For the second claim, since $T - sI$ is almost invertible, $(R_*)_{i, i} > 0$ for $i < n$. From the previous lemma,

$$(b \circ \Phi_*)(T, s) = \frac{(R_*)_{n, n}}{(R_*)_{n-1, n-1}} b(T);$$

if $b(T) = 0$ and $s = (T)_{n, n}$ then $(R_*)_{n, n} = 0$ and $b \circ \Phi_*$ is a product of two smooth functions, both zero, yielding $\text{grad}(b \circ \Phi_*) = 0$. \blacksquare

The operation of changing subdiagonal signs, i.e., of conjugation by some $E \in \mathcal{E}$, behaves well with respect to Φ and Φ_* . For $1 \leq j < n$, let $E_j \in \mathcal{E}$ be defined by

$$(E_j)_{i, i} = \begin{cases} 1, & i \leq j, \\ -1, & i > j. \end{cases}$$

Together with $-I$, the matrices E_i generate \mathcal{E} . For $T \in \mathcal{T}$, $\eta_i(T) = E_i T E_i$ differs from T only in the sign of the i -th subdiagonal coordinate: $(\eta_i(T))_{i+1, i} = -(T)_{i+1, i}$. The nontrivial involutions in \mathcal{T}_Λ are therefore generated by η_i , $1 \leq i < n$.

Lemma 2.4 *The domains $\text{Dom}(\Phi)$ and $\text{Dom}(\Phi_*)$ are \mathcal{E} -invariant and*

$$\Phi(\eta(T), s) = \eta(\Phi(T, s)), \quad \Phi_*(\eta(T), s) = \eta(\Phi_*(T, s)).$$

If $\det(T - sI) > 0$ then $\Phi(T, s) = \Phi_(T, s)$; if $\det(T - sI) < 0$, $\Phi(T, s) = \eta_{n-1}(\Phi_*(T, s))$; if $\det(T - sI) = 0$ and $(T, s) \in \text{Dom}(\Phi_*)$, then $b(\Phi_*(T, s)) = 0$.*

Proof: For $(T, s) \in \text{Dom}(\Phi)$, the matrices $T - sI$ and $E(T - sI)E$ are both invertible. The QR factorization $T - sI = QR$ yields $ETE - E(sI)E = (EQE)(ERE)$, preserving the positivity of the diagonal entries of the triangular part, so

$$\Phi(ETE, s) = (EQE)^* ETE(EQE) = EQ^* TQE = E\Phi(T, s)E.$$

The argument is similar for Φ_* . The claims for $T - sI$ invertible follow from the relation between Q and Q_* in Proposition 2.1; the case $\det(T - sI) = 0$ is a repetition of Lemma 2.3. \blacksquare

We are only interested in the case when the spectrum of T is simple, since a double eigenvalue implies reducibility. Since either version of shifted QR step preserves spectrum, restriction defines smooth maps $\Phi : (\mathcal{T}_\Lambda \times \mathbb{R}) \cap \text{Dom}(\Phi) \rightarrow \mathcal{T}_\Lambda$ and $\Phi_\star : (\mathcal{T}_\Lambda \times \mathbb{R}) \cap \text{Dom}(\Phi_\star) \rightarrow \mathcal{T}_\Lambda$.

Still in \mathcal{T}_Λ , it is convenient to consider the *step* $F_s(T) = \Phi_\star(T, s)$. For s not an eigenvalue of Λ , the domain of F_s is \mathcal{T}_Λ . The natural domain for F_{λ_i} instead is the *deflation domain* \mathcal{D}_Λ^i , the open dense subset of \mathcal{T}_Λ of matrices T for which $T - \lambda_i I$ is almost invertible. In other words, $T \in \mathcal{D}_\Lambda^i$ if and only if λ_i is an eigenvalue of the lowest irreducible block of T .

The definition of the step F_s differs from the usual one in that we use Φ_\star instead of Φ . Given Lemma 2.4, considerations about deflation are unaffected and our choice has the advantage of being smooth (and well defined) in \mathcal{D}_Λ^i .

The (i -th) *deflation set* is

$$\mathcal{D}_{\Lambda,0}^i = \{T \in \mathcal{T}_\Lambda \mid b(T) = 0, (T)_{n,n} = \lambda_i\}.$$

Since the spectrum of Λ is simple, $\mathcal{D}_{\Lambda,0}^i \subset \mathcal{D}_\Lambda^i$. Also, if $i \neq j$ then $\mathcal{D}_\Lambda^i \cap \mathcal{D}_{\Lambda,0}^j = \emptyset$. We saw in Lemma 2.3 that when the shift is taken to be an eigenvalue, a single step deflates a matrix, i.e., that the image of F_{λ_i} is contained in $\mathcal{D}_{\Lambda,0}^i$: we shall see in Proposition 2.5 that this image is in fact equal to $\mathcal{D}_{\Lambda,0}^i$.

Proposition 2.5 *If s is not an eigenvalue of Λ , the map $F_s : \mathcal{T}_\Lambda \rightarrow \mathcal{T}_\Lambda$ is a diffeomorphism. The image of $F_{\lambda_i} : \mathcal{D}_\Lambda^i \rightarrow \mathcal{T}_\Lambda$ is $\mathcal{D}_{\Lambda,0}^i$. The restriction $F_{\lambda_i}|_{\mathcal{D}_{\Lambda,0}^i} : \mathcal{D}_{\Lambda,0}^i \rightarrow \mathcal{D}_{\Lambda,0}^i$ is a diffeomorphism.*

Proof: If s is not an eigenvalue, compute $F_s^{-1}(T)$ by factoring $T - sI$ as RQ , R upper triangular with the first $n - 1$ diagonal entries positive and $Q \in SO(n)$: we claim that $F_s(T_0) = T$ for $T_0 = QR + sI$, proving that F_s is a diffeomorphism. Indeed, $QR = T_0 - sI$ is a $Q_\star R_\star$ factorization and thus $F_s(T_0) = Q_\star^* T_0 Q = T$.

From the last sentence of Section 2, the image of F_{λ_i} is contained in $\mathcal{D}_{\Lambda,0}^i \subset \mathcal{D}_\Lambda^i$. The fact that the restriction of F_{λ_i} to $\mathcal{D}_{\Lambda,0}^i$ is a diffeomorphism is proved as in the previous paragraph. ■

Commutativity of steps is well known and related to the complete integrability of the interpolating Toda flows ([6], [11], [13], [14]). For the reader's convenience we provide a quick proof.

Proposition 2.6 *Steps commute: $F_{s_0} \circ F_{s_1} = F_{s_1} \circ F_{s_0}$ in the appropriate domains.*

The domain of $F_{s_0} \circ F_{s_1} = F_{s_1} \circ F_{s_0}$ is \mathcal{T}_Λ if neither s_0 nor s_1 is an eigenvalue, \mathcal{D}_Λ^i if $s_0 = \lambda_i$ and s_1 is not an eigenvalue (or vice-versa) and the empty set in the rather pointless case $s_0 = \lambda_i, s_1 = \lambda_j, i \neq j$.

Proof: We prove commutativity only when s_0 and s_1 are not eigenvalues; the other cases follow easily. Consider $Q_\star R_\star$ factorizations

$$\begin{aligned} T - s_0 I &= Q_0 R_0, & T - s_1 I &= Q_1 R_1, \\ (T - s_0 I)(T - s_1 I) &= (T - s_1 I)(T - s_0 I) = Q_2 R_2. \end{aligned}$$

For $F_{s_0}(T) - s_1 = Q_0^*(T - s_1)Q_0 = Q_3 R_3$, we have $F_{s_1}(F_{s_0}(T)) = Q_3^* F_{s_0}(T) Q_3 = Q_3^* Q_0^* T Q_0 Q_3$. Thus

$$Q_0^*(T - s_1)Q_0 R_0 = Q_0^*(T - s_1 I)(T - s_0 I) = Q_0^* Q_2 R_2 = Q_3 R_3 R_0$$

and therefore $Q_0^* Q_2 = Q_3$ and $F_{s_0}(F_{s_1}(T)) = Q_2^* T Q_2$. ■

3 Simple shift strategies

The point of using a shift strategy is to accelerate deflation, ideally by choosing s near an eigenvalue of T . A *simple shift strategy* is an \mathcal{E} -invariant function $\sigma : \mathcal{T}_\Lambda \rightarrow \mathbb{R}$ such that there exists $C_\sigma > 0$ such that for all $T \in \mathcal{T}_\Lambda$ there is an eigenvalue λ_i with $|\sigma(T) - \lambda_i| \leq C_\sigma |b(T)|$. In particular, if $T \in \mathcal{D}_{\Lambda,0}^i$ then $\sigma(T) = \lambda_i$.

The *step* associated with a (simple) shift strategy σ is F_σ , defined by $F_\sigma(T) = F_{\sigma(T)}(T)$. The natural domain for F_σ is the set of matrices T for which $T - \sigma(T)I$ is almost invertible. From Section 2, it includes all unreduced matrices and open neighborhoods of each deflation set $\mathcal{D}_{\Lambda,0}^i$. We shall also see in Section 6 that it contains a dense open subset $\mathcal{U}_{\Lambda,\epsilon}$ of \mathcal{T}_Λ invariant under F_σ . A more careful description of this domain will not be needed.

Quoting Parlett ([14]), there are shifts for all seasons. Let ρ be *Rayleigh's shift*: $\rho(T) = (T)_{n,n}$. Denote the bottom 2×2 diagonal principal minor of a matrix $T \in \mathcal{T}$ by \hat{T} : *Wilkinson's shift* $\omega(T)$ is the eigenvalue of \hat{T} closer to $(T)_{n,n}$ (in case of draw, take the smallest eigenvalue).

Lemma 3.1 *The functions ρ and ω are simple shift strategies with $C_\rho = \sqrt{2}$ and $C_\omega = 2\sqrt{2}$.*

We use here the Wielandt-Hoffman theorem (for a simple proof using the Toda dynamics, see [4]): if $S, T \in \mathcal{T}$ have eigenvalues σ_i and λ_i in increasing order then

$$\sum_i |\sigma_i - \lambda_i|^2 \leq \text{tr}((S - T)^2).$$

Proof: Invariance is trivial for ρ ; for ω , it follows from the fact that changing signs of off-diagonal entries of a 2×2 matrix does not change its spectrum.

Let $B = e_n e_{n-1}^* + e_{n-1} e_n^*$ and $S = T - b(T)B$ so that $\rho(T) = (T)_{n,n}$ is an eigenvalue of S . From the Wielandt-Hoffman theorem, for some i ,

$$|\rho(T) - \lambda_i| \leq \sqrt{2} |b(T)|,$$

proving that $C_\rho = \sqrt{2}$. Apply again the Wielandt-Hoffman theorem to the 2×2 trailing principal minors of S and T to deduce that

$$|(T)_{n,n} - \omega(T)| \leq \sqrt{2} |b(T)|.$$

We thus have $|\omega(T) - \lambda_i| \leq 2\sqrt{2} |b(T)|$ and $C_\omega = 2\sqrt{2}$, as desired. \blacksquare

Another example of (simple) shift strategy, the *mixed Wilkinson-Rayleigh strategy*, uses Wilkinson's shift unless the matrix is already near deflation, in which case we use Rayleigh's:

$$\sigma(T) = \begin{cases} \rho(T), & |(T)_{n,n-1}| < \epsilon, \\ \omega(T), & |(T)_{n,n-1}| \geq \epsilon; \end{cases}$$

here $\epsilon > 0$ is a small constant.

Simple shift strategies are not required to be continuous and ω is definitely not. For a simple shift strategy σ , let $\mathcal{S}_\sigma \subset \mathcal{T}_\Lambda$ be the *singular support* of σ , i.e., a minimal closed set on whose complement σ is smooth. For example, \mathcal{S}_ω is the set of matrices $T \in \mathcal{T}_\Lambda$ for which the two eigenvalues $\omega_-(T)$ and $\omega_+(T)$ of \hat{T} are equidistant from $(T)_{n,n}$, or, equivalently, for which $(T)_{n,n} = (T)_{n-1,n-1}$. The set \mathcal{S}_σ will play an important role later.

We consider the phase portrait of F_ω for 3×3 matrices. In this case, the reader may check that the domain of F_ω is the full set \mathcal{T}_Λ . Let $\mathcal{J}_\Lambda \subset \mathcal{T}_\Lambda$ be set of *Jacobi matrices* similar to Λ , i.e., matrices $T \in \mathcal{T}_\Lambda$ with strictly positive subdiagonal entries.

Recall that the closure $\tilde{\mathcal{J}}_\Lambda \subset \mathcal{T}_\Lambda$ is diffeomorphic to a hexagon, the permutohedron in this dimension. The set $\tilde{\mathcal{J}}_\Lambda$ is not invariant under F_ω but we may define $\tilde{F}_\omega(T)$ with $\tilde{F}_\omega : \tilde{\mathcal{J}}_\Lambda \rightarrow \tilde{\mathcal{J}}_\Lambda$ by dropping signs of subdiagonal entries of $F_\omega(T)$. As discussed above, this standard procedure is mostly harmless.

Two examples of \tilde{F}_ω are given in Figure 3, which represent $\tilde{\mathcal{J}}_\Lambda$ for the $\Lambda = \text{diag}(1, 2, 4)$ on the left and $\Lambda = \text{diag}(-1, 0, 1)$ on the right. The vertices are the six diagonal matrices similar to Λ and the edges consist of reduced matrices. Labels indicate the diagonal entries of the corresponding matrices. Three edges form $\mathcal{D}_{\Lambda,0}^i \cap \tilde{\mathcal{J}}_\Lambda$: they alternate, starting from the bottom horizontal edge on both hexagons. The set $\mathcal{S} \cap \tilde{\mathcal{J}}_\Lambda$ is indicated in both cases.




Figure 3: The phase space of Wilkinson's step for $n = 3$.

Vertices are fixed points of \tilde{F}_ω and boundary edges are invariant sets. A simple arrow indicates the motion of the points $\tilde{F}_\omega^k(T)$ along the edge. Points T on an arc with a double arrow are taken to a diagonal matrix in a single step: the arc points to $\tilde{F}_\omega(T)$. Arcs marked with a transversal segment consist of fixed points of \tilde{F}_ω .

Points on both sides of \mathcal{S}_ω are taken far apart: there is a jump discontinuity along \mathcal{S}_ω . From Theorem 2, the decay of the bottom subdiagonal entry under Wilkinson's step away from $\mathcal{S}_\omega \cap \mathcal{D}_{\Lambda,0}$ is cubic. As discussed in [9], near $\mathcal{S}_\omega \cap \mathcal{D}_{\Lambda,0}$ this decay is quadratic, but not cubic. For the left hexagon, cubic convergence occurs in the long run because the sequence $\tilde{F}_\omega^k(T)$ stays close to this intersection only for a few values of k , illustrating Theorem 3.

In the case $\Lambda = \text{diag}(-1, 0, 1)$, the bottom edge consists of fixed points. As mentioned in the Introduction (see Figure 2), this case has a special asymptotic behavior: the (fixed) point labeled by $(0, 0, 0)$ is the central point of the set \mathcal{X} . If $T \in \mathcal{X}$ then the sequence $(\tilde{F}_\omega^k(T))$ is contained in \mathcal{X} and converges to the central point at a strictly quadratic rate.

4 Tubular coordinates

Recall that a map $\Pi : X \rightarrow Y \subset X$ is a *projection* if $\Pi(X) = Y$ and $\Pi \circ \Pi = \Pi$. Instead of using abstract topological facts to prove the existence of some projection $\mathcal{D}_\Lambda^i \rightarrow \mathcal{D}_{\Lambda,0}^i$ we prefer to construct a specific projection which works well with the QR steps. The map $F_{\lambda_i} : \mathcal{D}_\Lambda^i \rightarrow \mathcal{D}_{\Lambda,0}^i$ is not a projection but can, using Proposition 2.5, be used to define one: the *canonical projection* $\Pi_i : \mathcal{D}_\Lambda^i \rightarrow \mathcal{D}_{\Lambda,0}^i$,

$$\Pi_i(T) = (F_{\lambda_i}|_{\mathcal{D}_{\Lambda,0}^i})^{-1}(F_{\lambda_i}(T)).$$

Proposition 4.1 *The map Π_i is a smooth projection which commutes with steps: $\Pi_i(F_s(T)) = F_s(\Pi_i(T))$ provided s is not an eigenvalue of Λ different from λ_i .*

Proof: The map Π_i is clearly smooth and, for $T \in \mathcal{D}_{\Lambda,0}^i$, we have

$$\Pi_i(T) = (F_{\lambda_i}|_{\mathcal{D}_{\Lambda,0}^i})^{-1}(F_{\lambda_i}(T)) = T,$$

proving that Π_i is a projection. Commutativity follows from Proposition 2.6. \blacksquare

For a diagonal matrix Λ with simple spectrum and $\epsilon > 0$, the *deflation neighborhood* $\mathcal{D}_{\Lambda,\epsilon} \subset \mathcal{T}_\Lambda$ is the closed set of matrices $T \in \mathcal{T}_\Lambda$ with $|b(T)| \leq \epsilon$. This notation is consistent with $\mathcal{D}_{\Lambda,0}$ for the deflation set. As we shall see in Propositions 4.2 and 5.1, for sufficiently small $\epsilon > 0$ the set $\mathcal{D}_{\Lambda,\epsilon}$ has connected components $\mathcal{D}_{\Lambda,\epsilon}^i \subset \mathcal{D}_\Lambda^i$, $\mathcal{D}_{\Lambda,\epsilon}^i \supset \mathcal{D}_{\Lambda,0}^i$, which are invariant under steps F_s for shifts s near λ_i , i.e., $F_s(\mathcal{D}_{\Lambda,\epsilon}^i) \subset \mathcal{D}_{\Lambda,\epsilon}^i$. The sets $\mathcal{D}_{\Lambda,\epsilon}^i$ are therefore also invariant under F_σ .

Denote the distance between a matrix T and a compact set of matrices \mathcal{N} by $\text{dist}(T, \mathcal{N}) = \min_{S \in \mathcal{N}} \|T - S\|$. Let $\gamma = \min_{i \neq j} |\lambda_i - \lambda_j|$ be the *spectral gap* of Λ and $B = e_n e_{n-1}^* + e_{n-1} e_n^*$.

Recall that if \mathcal{N} is a submanifold of codimension k of \mathcal{M} then a *closed tubular neighborhood* of \mathcal{N} consists of a closed neighborhood \mathcal{N}_ϵ of \mathcal{N} and a diffeomorphism $\zeta : \mathcal{N}_\epsilon \rightarrow \mathcal{N} \times \mathbb{B}_\epsilon^k$ with $\zeta(x) = (x, 0)$ for $x \in \mathcal{N}$ (here $\mathbb{B}_\epsilon^k \subset \mathbb{R}^k$ is the closed ball of radius ϵ around the origin). Given $x \in \mathcal{N}$, the preimage $\zeta^{-1}(\{x\} \times \mathbb{B}_\epsilon^k)$ is a manifold with boundary of dimension k , the *fiber* through x . We now construct tubular neighborhoods of the deflation sets $\mathcal{D}_{\Lambda,0}^i$; here the codimension is $k = 1$.

Proposition 4.2 *Each $\mathcal{D}_{\Lambda,0}^i \subset \mathcal{T}_\Lambda$ is a compact submanifold of codimension 1 diffeomorphic to \mathcal{T}_{Λ_i} , where $\Lambda_i = \text{diag}(\lambda_1, \dots, \lambda_{i-1}, \lambda_{i+1}, \dots, \lambda_n)$. There exists $\epsilon_{\text{tub}} > 0$ such that for $\epsilon \in (0, \epsilon_{\text{tub}})$:*

- (a) *the connected components $\mathcal{D}_{\Lambda,\epsilon}^i$ of $\mathcal{D}_{\Lambda,\epsilon}$ consist of matrices $T \in \mathcal{D}_{\Lambda,\epsilon}$ for which $|(T)_{n,n} - \lambda_i| < \sqrt{2}\epsilon$;*
- (b) *the map $\zeta : \mathcal{D}_{\Lambda,\epsilon}^i \rightarrow \mathcal{D}_{\Lambda,0}^i \times [-\epsilon, \epsilon]$ given by $\zeta(T) = (\Pi_i(T), b(T))$ is a closed tubular neighborhood of $\mathcal{D}_{\Lambda,0}^i$;*
- (c) *there is a constant $C_b > 0$ such that for all $T \in \mathcal{D}_{\Lambda,\epsilon}^i$,*

$$|b(T)| \leq \text{dist}(T, \mathcal{D}_{\Lambda,0}^i) \leq \|T - \Pi_i(T)\| \leq C_b |b(T)|.$$

Proof: We first show that the gradient of the restriction $b|_{\mathcal{T}_\Lambda}$ at a point $T_{\mathcal{D}} \in \mathcal{D}_{\Lambda,0}$ is not zero. Consider the characteristic polynomial along the line $T_{\mathcal{D}} + tB$: this is a smooth even function of t and therefore B is tangent to \mathcal{T}_Λ at $T_{\mathcal{D}}$, the point on which $t = 0$. On the other hand, the directional derivative of b along the same line equals 1. Thus $\mathcal{D}_{\Lambda,0} \subset \mathcal{T}_\Lambda$ is a submanifold of codimension 1. The diffeomorphism with \mathcal{T}_{Λ_i} takes T to \hat{T} , the leading $(n-1) \times (n-1)$ principal minor of T .

Assume $\epsilon < \gamma/(2\sqrt{2})$. Consider matrices $T \in \mathcal{D}_{\Lambda,\epsilon}$ and $S = T - b(T)B$, so that $(T)_{n,n}$ is an eigenvalue of S . By the Wielandt-Hoffman theorem, there exists an index i for which $|(T)_{n,n} - \lambda_i| < \sqrt{2}\epsilon$, defining the sets $\mathcal{D}_{\Lambda,\epsilon}^i$ (at this point we do not yet know that $\mathcal{D}_{\Lambda,\epsilon}^i$ is connected).

For $T_{\mathcal{D}} \in \mathcal{D}_{\Lambda,0}^i$, the derivative $D\Pi_i(T_{\mathcal{D}})$ equals the identity on the subspace tangent to $\mathcal{D}_{\Lambda,0}^i$ and has a kernel of dimension 1. Thus, for sufficiently small ϵ_{tub} , item (b) holds. This also proves that each $\mathcal{D}_{\Lambda,\epsilon}^i$ is connected, completing the proof of item (a).

The first two inequalities in (c) are trivial. Now

$$\|T - \Pi_i(T)\| = \|\zeta^{-1}(\Pi_i(T), b(T)) - \zeta^{-1}(\Pi_i(T), 0)\| \leq C_b |b(T)|,$$

where the derivative of $\zeta^{-1}(T_{\mathcal{D}}, \delta)$ with respect to the second coordinate is bounded by C_b on the compact set $\mathcal{D}_{\Lambda,0} \times [-\epsilon_{\text{tub}}, \epsilon_{\text{tub}}]$. \blacksquare

The diffeomorphism ζ defines *tubular coordinates* for $T \in \mathcal{D}_{\Lambda,\epsilon}^i$: the matrix $\Pi_i(T) \in \mathcal{D}_{\Lambda,0}^i \approx \mathcal{T}_{\Lambda_i}$ and $b(T)$. Under tubular coordinates, QR steps with shift are given by a simple formula.

Corollary 4.3 Consider Λ , i and $\epsilon \in (0, \epsilon_{\text{tub}})$. Then

$$\zeta \circ F_s \circ \zeta^{-1} : \mathcal{D}_{\Lambda,0}^i \times [-\epsilon, \epsilon] \rightarrow \mathcal{D}_{\Lambda,0}^i \times [-\epsilon, \epsilon]$$

$$(T, b) \mapsto \left(F_s(T), \frac{(R_\star)_{n,n}}{(R_\star)_{n-1,n-1}} b \right)$$

where $\zeta^{-1}(T, b) - sI = Q_\star R_\star$.

Proof: This follows directly from Lemma 2.2 and Propositions 4.1 and 4.2. \blacksquare

5 Convergence to deflation

Sufficiently thin deflation neighborhoods $\mathcal{D}_{\Lambda,\epsilon}^i$ are invariant under F_s for $s \approx \lambda_i$.

Proposition 5.1 Given $C > 0$, there exists $\epsilon_{\text{inv}} \in (0, \epsilon_{\text{tub}})$, such that for any $\epsilon \in (0, \epsilon_{\text{inv}})$ and $s \in [\lambda_i - C\epsilon, \lambda_i + C\epsilon]$ we have $F_s(\mathcal{D}_{\Lambda,\epsilon}^i) \subset \text{int}(\mathcal{D}_{\Lambda,\epsilon/2}^i)$.

For a simple shift strategy $\sigma : \mathcal{T}_\Lambda \rightarrow \mathbb{R}$, there exists $\epsilon_{\text{inv}} > 0$ such that if $\epsilon \in (0, \epsilon_{\text{inv}})$ then $F_\sigma(\mathcal{D}_{\Lambda,\epsilon}^i) \subset \text{int}(\mathcal{D}_{\Lambda,\epsilon/2}^i)$.

In particular, F_s is well defined in $\mathcal{D}_{\Lambda,\epsilon}^i$ for $\epsilon \in (0, \epsilon_{\text{inv}})$.

Proof: Recall that $F_s(\mathcal{D}_{\Lambda,0}^i) = \mathcal{D}_{\Lambda,0}^i$. From Lemma 2.3, the derivative of $b \circ \Phi_\star$ is zero at $\mathcal{D}_{\Lambda,0}^i \times \{\lambda_i\}$. Compactness of $\mathcal{D}_{\Lambda,0}^i$ thus implies that in a sufficiently small neighborhood of $\mathcal{D}_{\Lambda,0}^i \times \{\lambda_i\}$ we have $|b(F_s(T))| \leq |b(T)|/3$.

Now consider a simple shift strategy σ : there exists $C_\sigma > 0$ such that $|\sigma(T) - \lambda_i| < C_\sigma b(T)$; apply the first statement with $C = C_\sigma$. \blacksquare

Thus, F_σ squeezes neighborhoods $\mathcal{D}_{\Lambda,\epsilon}^i$ at least linearly. Equivariance and smoothness imply an estimate stronger than that in the definition of simple shift strategy. We do not want to assume, however, that $\mathcal{D}_{\Lambda,0} \cap \mathcal{S}_\sigma = \emptyset$: after all, this is not true even for Wilkinson's shift. We need a more careful statement.

Lemma 5.2 Consider a shift strategy σ and ϵ_{inv} as in Proposition 5.1. For a compact set $\mathcal{K} \subset \mathcal{D}_{\Lambda,\epsilon_{\text{inv}}}^i \setminus (\mathcal{D}_{\Lambda,0}^i \cap \mathcal{S}_\sigma)$, there exists $C_{\mathcal{K}}$ such that for all $T \in \mathcal{K}$ we have $|\sigma(T) - \lambda_i| \leq C_{\mathcal{K}} b(T)^2$.

Proof: Let $\mathcal{K}_{\mathcal{D}} = \mathcal{K} \cap \mathcal{D}_{\Lambda,0}^i$; enlarge $\mathcal{K}_{\mathcal{D}}$ along $\mathcal{D}_{\Lambda,0}^i$ to obtain another compact set $\mathcal{K}_1 \subset \mathcal{D}_{\Lambda,0}^i \setminus \mathcal{S}_\sigma$, $\mathcal{K}_{\mathcal{D}} \subset \text{int}_{\mathcal{D}_{\Lambda,0}^i}(\mathcal{K}_1)$. Fatten \mathcal{K}_1 along fibers to define $\tilde{\mathcal{K}}_1 = \zeta^{-1}(\mathcal{K}_1 \times [-\epsilon, \epsilon])$, $\epsilon \in (0, \epsilon_{\text{inv}})$, which, without loss, still avoids \mathcal{S}_σ . For each $T_{\mathcal{D}} \in \mathcal{K}_1$, consider the function $h_{T_{\mathcal{D}}}(b) = \sigma(\zeta^{-1}(T_{\mathcal{D}}, b))$, obtained by restricting σ to a fiber of $\mathcal{D}_{\Lambda,\epsilon}^i$. Each $h_{T_{\mathcal{D}}}$ is smooth and even and therefore satisfies $|h_{T_{\mathcal{D}}}(b) - \lambda_i| \leq C_{T_{\mathcal{D}}} |b|^2$. By compactness, there exists $C_{\mathcal{K}_1}$ such that $|h_{T_{\mathcal{D}}}(b) - \lambda_i| \leq C_{\mathcal{K}_1} |b|^2$ for all $T_{\mathcal{D}} \in \mathcal{K}_1$. In other words, there exists $C_{\tilde{\mathcal{K}}_1}$ such that $|\sigma(T) - \lambda_i| \leq C_{\tilde{\mathcal{K}}_1} |b(T)|^2$ for all $T \in \tilde{\mathcal{K}}_1$. The estimate for $T \notin \tilde{\mathcal{K}}_1$ is trivial. \blacksquare

Proof of Theorem 2: Take $\epsilon = \epsilon_{\text{inv}}$ as in Proposition 5.1 so that $\mathcal{D}_{\Lambda,\epsilon}^i$ is invariant under F_σ .

Let $\varphi = b \circ \Phi_\star$. We compute the Taylor expansion of $\varphi(T, s)$ at $(T_{\mathcal{D}}, \lambda_i)$, $T_{\mathcal{D}} \in \mathcal{D}_{\Lambda,0}^i$: from Lemma 2.3, the gradient of φ at $(T_{\mathcal{D}}, \lambda_i)$ is zero. Thus, up to a third order remainder,

$$\begin{aligned} \varphi(T, s) &= \varphi(T_{\mathcal{D}}, \lambda_i) + \frac{1}{2} \varphi_{T,T}(T_{\mathcal{D}}, \lambda_i)(T - T_{\mathcal{D}}, T - T_{\mathcal{D}}) + \\ &\quad + \varphi_{T,s}(T_{\mathcal{D}}, \lambda_i)(T - T_{\mathcal{D}}, s - \lambda_i) + \frac{1}{2} \varphi_{s,s}(T_{\mathcal{D}}, \lambda_i)(s - \lambda_i, s - \lambda_i) + \\ &\quad + \text{Rem}_3(T - T_{\mathcal{D}}, s - \lambda_i). \end{aligned}$$

Now, $\varphi(T_{\mathcal{D}}, \lambda_i) = 0$ and, again from Lemma 2.3, $\varphi(T, \lambda_i) = 0$ for all $T \in \mathcal{T}_{\Lambda}$, hence $\varphi_{T,T}(T_{\mathcal{D}}, \lambda_i) = 0$. Let C_{σ} be the constant in the definition of a simple shift strategy. By compactness, there exists $C_1 > 0$ such that for all $T_{\mathcal{D}} \in \mathcal{D}_{\Lambda,0}^i$, $T \in \mathcal{D}_{\Lambda,\epsilon}^i$ and $s \in [\lambda_i - C_{\sigma}\epsilon, \lambda_i + C_{\sigma}\epsilon]$, we have

$$|\varphi(T, s)| \leq C_1 |s - \lambda_i| (\|T - T_{\mathcal{D}}\| + |s - \lambda_i|)$$

We now apply this estimate for $T_{\mathcal{D}} = \Pi_i(T)$, where $T \in \mathcal{D}_{\Lambda,\epsilon}^i$. By Proposition 4.2, since $\epsilon < \epsilon_{\text{tub}}$, $\|T - T_{\mathcal{D}}\| = \|T - \Pi_i(T)\| \leq C_b |b(T)|$ and therefore

$$|\varphi(T, s)| \leq C_1 |s - \lambda_i| (C_b |b(T)| + |s - \lambda_i|)$$

implying the quadratic estimate

$$|b(F_{\sigma}(T))| = |\varphi(T, \sigma(T))| \leq C_1 |\sigma(T) - \lambda_i| (C_b |b(T)| + |\sigma(T) - \lambda_i|) \leq C_q |b(T)|^2.$$

Using Lemma 5.2 yields the cubic estimate in (c). \blacksquare

As a corollary, we obtain the well known fact that, near deflation, the rate of convergence of Rayleigh's (as well as the mixed Wilkinson-Rayleigh) strategy has cubic convergence. The rate of convergence for Wilkinson's strategy is subtler.

We construct a larger invariant set for F_{σ} . Let $\mathcal{U}_{\Lambda} \subset \mathcal{T}_{\Lambda}$ be the set of unreduced matrices; for $\epsilon > 0$, let $\mathcal{U}_{\Lambda,\epsilon} = \mathcal{U}_{\Lambda} \cup \text{int}(\mathcal{D}_{\Lambda,\epsilon})$. Notice that $\mathcal{U}_{\Lambda,\epsilon}$ is open, dense and path-connected.

Lemma 5.3 *For a shift strategy $\sigma : \mathcal{T}_{\Lambda} \rightarrow \mathbb{R}$, ϵ_{inv} as in Proposition 5.1 and $\epsilon \in (0, \epsilon_{\text{inv}})$, the open set $\mathcal{U}_{\Lambda,\epsilon}$ is invariant under F_{σ} .*

Proof: If $T \in \mathcal{U}_{\Lambda}$ and $\sigma(T)$ is not in the spectrum then $F_{\sigma}(T)$ is (well defined and) unreduced. If $T \in \mathcal{U}_{\Lambda}$ and $\sigma(T) = \lambda_i$ then $F_{\sigma}(T) \in \mathcal{D}_{\Lambda,0}^i \subset \mathcal{U}_{\Lambda,\epsilon}$. Finally, if $T \in \text{int}(\mathcal{D}_{\Lambda,\epsilon}^i)$ then, by Proposition 5.1, $F_{\sigma}(T) \in \text{int}(\mathcal{D}_{\Lambda,\epsilon/2}^i) \subset \mathcal{U}_{\Lambda,\epsilon}$. \blacksquare

Notice that we do not assume σ or F_{σ} to be continuous. This shows that for F_{σ} defined from a simple shift strategy σ the extra hypothesis in Theorem 1, item (b), actually holds: just take $\mathcal{K}_i = \mathcal{D}_{\Lambda,\epsilon}^i$.

A simple shift strategy σ is *deflationary* if for any $T \in \mathcal{U}_{\Lambda,\epsilon_{\text{inv}}}$ there exists $K \in \mathbb{N}$ such that $F_{\sigma}^K(T) \in \mathcal{D}_{\Lambda,\epsilon_{\text{inv}}}$. It is now a corollary of Theorem 1 and Lemma 5.3 that continuous simple shift strategies are not deflationary.

Rayleigh's strategy is known not to be deflationary. The following well known estimate ([7] and [14], section 8-10) implies that Wilkinson's strategy is not only deflationary but uniformly so, in the sense that there exists K with $F_{\omega}^K(\mathcal{U}_{\Lambda,\epsilon_{\text{inv}}}) \subset \mathcal{D}_{\Lambda,\epsilon_{\text{inv}}}$. As a corollary, the mixed Wilkinson-Rayleigh strategy is also uniformly deflationary provided $\epsilon > 0$ is sufficiently small.

Fact 5.4 *For $T \in \mathcal{T}$ and $k \in \mathbb{N}$,*

$$|b(F_{\omega}^k(T))|^3 \leq \frac{|b(T)^2(T)_{n-1,n-2}|}{(\sqrt{2})^{k-1}}.$$

In [14], the result is shown for unreduced matrices; the case $T \in \mathcal{U}_{\Lambda,\epsilon_{\text{inv}}}$ follows by taking limits. Notice that for $T \in \mathcal{T}_{\Lambda}$, the numerator $|b(T)^2(T)_{n-1,n-2}|$ is uniformly bounded.

6 Dynamics for a.p. free spectra

From the previous section, cubic convergence may be lost when the orbit $F_\sigma^k(T)$ passes near the set $\mathcal{S}_\sigma \cap \mathcal{D}_{\Lambda,0}$. Our next task is to measure when this happens, by studying the dynamics associated to a shift strategy in a deflation neighborhood, i.e., the iterates of $F_\sigma : \mathcal{D}_{\Lambda,\epsilon}^i \rightarrow \mathcal{D}_{\Lambda,\epsilon}^i$, $\epsilon \in (0, \epsilon_{\text{inv}})$. Most of what we need can be read in the projection onto $\mathcal{D}_{\Lambda,0}^i$, where F_σ coincides with F_{λ_i} .

A matrix $T \in \mathcal{T}$ with simple spectrum is *a.p. free* if no three eigenvalues are in arithmetic progression and a.p. otherwise. Different kinds of spectra lead to different dynamics: in this section we handle the a.p. free case, clearly a generic restriction. Let \tilde{T} be the leading principal $(n-1) \times (n-1)$ minor of T . The following result is standard.

Proposition 6.1 *Let $\Lambda \in \mathcal{T}$ be an $n \times n$ diagonal a.p. free matrix with spectrum $\lambda_1 < \dots < \lambda_n$. For each i , consider $F_{\lambda_i} : \mathcal{D}_{\Lambda,0}^i \rightarrow \mathcal{D}_{\Lambda,0}^i$ as above. For any $T \in \mathcal{D}_{\Lambda,0}^i$, the sequence $(F_{\lambda_i}^k(T))$ converges to a diagonal matrix.*

Proof: The map F_{λ_i} on $\mathcal{D}_{\Lambda,0}^i$ amounts to a *QR* step with shift λ_i on \tilde{T} , which has eigenvalues λ_j , $j \neq i$. The a.p. free hypothesis implies that the absolute values of the eigenvalues of $\tilde{T} - \lambda_i I$ are distinct. If \tilde{T} is unreduced then, as is well known, the standard *QR* iteration converges to a diagonal matrix, with diagonal entries in decreasing order of absolute value. More generally, if \tilde{T} is reduced, apply the above result to each unreduced sub-block. ■

We shall use *height functions* for the *QR* steps F_s , s near λ_i , i.e., functions $H_i : \mathcal{D}_{\Lambda,\epsilon}^i \rightarrow \mathbb{R}$ with $H_i(F_s(T)) > H_i(T)$ provided T is not diagonal. Such height functions and related scenarios have been considered in [1], [4], [11] and [16].

The matrix $W = \text{diag}(w_1, \dots, w_n)$ is a *weight matrix* if $w_1 > \dots > w_n$. Since Λ is a.p. free, there exists $\epsilon_{\text{ap}} \in (0, \epsilon_{\text{inv}})$ such that if $s \in \mathcal{I}_i = [\lambda_i - \epsilon_{\text{ap}}, \lambda_i + \epsilon_{\text{ap}}]$ then the numbers $|\lambda_j - s|$ are distinct and their order does not depend on s .

Proposition 6.2 *Let Λ be an a.p. free diagonal matrix, W a weight matrix and ϵ_{ap} as above. For $\delta_H > 0$, set $L_i(x) = \log((x - \lambda_i)^2 + \delta_H)$ and let $H_i : \mathcal{D}_{\Lambda,\epsilon_{\text{ap}}}^i \rightarrow \mathbb{R}$ be defined by $H_i(T) = \text{tr}(WL_i(T))$. There exists $\delta_H > 0$ such that*

$$\max_{T \in \partial \mathcal{D}_{\Lambda,\epsilon_{\text{ap}}}^i} H_i(T) < \min_{T \in \mathcal{D}_{\Lambda,0}^i} H_i(T)$$

and, for any $s \in \mathcal{I}_i$, H_i is a height function for $F_s : \mathcal{D}_{\Lambda,\epsilon_{\text{ap}}}^i \rightarrow \mathcal{D}_{\Lambda,\epsilon_{\text{ap}}}^i$.

Here, $L_i(T) = X \text{diag}(L_i(\lambda_1), \dots, L_i(\lambda_n)) X^{-1}$ for $T = X \Lambda X^{-1}$ so that if p is a polynomial and $L_i(\lambda_j) = p(\lambda_j)$ for $j = 1, \dots, n$ then $L_i(T) = p(T)$. The only conditions on L_i which will be used in the proof are that $|\lambda_j - \lambda_i| < |\lambda_k - \lambda_i|$ implies $L_i(\lambda_j) < L_i(\lambda_k)$ and that $L_i(\lambda_i)$ is very negative (for small δ_H).

The proof requires some basic facts about *f-Q_{*}R_{*} steps* (again related to the integrability of the Toda lattice); these facts will not be used elsewhere. For a real diagonal matrix Λ with simple spectrum, let \mathcal{O}_Λ be the set of all real symmetric matrices similar to Λ ; it is well known that \mathcal{O}_Λ is a smooth compact manifold. The *f-Q_{*}R_{*} step* applied to a matrix $S \in \mathcal{O}_\Lambda$ is the map $F_f : \mathcal{A}_{\Lambda,f} \rightarrow \mathcal{O}_\Lambda$ defined by $F_f(S) = Q_*^* S Q_*$, where Q_* is obtained from the factorization $f(S) = Q_* R_*$ and $S \in \mathcal{A}_{\Lambda,f}$ if and only if $f(S)$ is almost invertible. If $T \in \mathcal{T}_\Lambda \cap \mathcal{A}_{\Lambda,f}$ then $F_f(T) \in \mathcal{T}_\Lambda$ (use the same proof as in Lemma 2.2). The maps $F_s : \mathcal{T}_\Lambda \rightarrow \mathcal{T}_\Lambda$ defined above correspond to restrictions of F_f for $f(x) = x - s$.

For a continuous function $h : \mathbb{R} \rightarrow \mathbb{R}$, if $S \in \mathcal{O}_\Lambda$ then the matrix function $h(S)$ belongs to \mathcal{O}_M , where $M = h(\Lambda)$. With the obvious abuse of notation, we have a diffeomorphism $h : \mathcal{O}_\Lambda \rightarrow \mathcal{O}_M$ provided h is injective in the spectrum of Λ .

Lemma 6.3 For h injective in the spectrum of Λ , consider the diffeomorphism $h : \mathcal{O}_\Lambda \rightarrow \mathcal{O}_M$, where $M = h(\Lambda)$. Let f and \tilde{f} be continuous functions defined in neighborhoods of the spectra of Λ and M , respectively, satisfying $\tilde{f}(h(\lambda_j)) = f(\lambda_j)$ for each j with QR steps $F_f : \mathcal{O}_\Lambda \rightarrow \mathcal{O}_\Lambda$ and $F_{\tilde{f}} : \mathcal{O}_M \rightarrow \mathcal{O}_M$. Then $h \circ F_f = F_{\tilde{f}} \circ h$.

Proof: The hypothesis implies that, for $T \in \mathcal{O}_\Lambda$, $f(T) = \tilde{f}(h(T)) = QR$ and hence $F_f(T) = Q^*TQ$ and $F_{\tilde{f}}(h(T)) = Q^*h(T)Q$. Thus $h(F_f(T)) = F_{\tilde{f}}(h(T))$. ■

Let I_r be the $n \times n$ truncated identity matrix, i.e., $(I_r)_{i,i} = 1$ for $i \leq r$, other entries being equal to zero.

Lemma 6.4 Let M be a diagonal matrix with simple spectrum and $\tilde{f} : \mathbb{R} \rightarrow \mathbb{R}$ be a function for which $\mu_i < \mu_j$ implies $|\tilde{f}(\mu_i)| < |\tilde{f}(\mu_j)|$. Consider the \tilde{f} -QR step $F_{\tilde{f}} : \mathcal{A}_{M,\tilde{f}} \rightarrow \mathcal{O}_M$. For any $S \in \mathcal{A}_{M,\tilde{f}}$ and $r = 1, \dots, n-1$, $\text{tr}(I_r F_{\tilde{f}}(S)) \geq \text{tr}(I_r S)$. For $r = 1$, equality only holds if $(S)_{1,j} = 0$ for all $j > 1$.

This argument follows closely the first proof in [4].

Proof: Let V_r be the range of I_r and $\mu_{r,j}(S)$ be the eigenvalues of the leading principal $r \times r$ minor of S , listed in nondecreasing order. We claim that $\mu_{r,j}(F_{\tilde{f}}(S)) \geq \mu_{r,j}(S)$, which immediately implies $\text{tr}(I_r F_{\tilde{f}}(S)) \geq \text{tr}(I_r S)$. Recall that $F_{\tilde{f}}(S) = Q_\star^* S Q_\star$ where $Q_\star R_\star = \tilde{f}(S)$. Let U be an upper triangular matrix such that $Q_\star u = \tilde{f}(S) U u$ for $u \in V_r$. By min-max,

$$\begin{aligned} \mu_{r,j}(S) &= \max_{A \subset V_r} \min_{u \in A \setminus \{0\}} \frac{\langle u, Su \rangle}{\langle u, u \rangle}, \\ &\quad \dim(A) = r+1-j \\ \mu_{r,j}(F_{\tilde{f}}(S)) &= \max_A \min_u \frac{\langle u, F_{\tilde{f}}(S)u \rangle}{\langle u, u \rangle} = \max_A \min_u \frac{\langle \tilde{f}(S)Uu, S\tilde{f}(S)Uu \rangle}{\langle \tilde{f}(S)Uu, \tilde{f}(S)Uu \rangle} \\ &= \max_{A'=UA} \min_{u' \in A' \setminus \{0\}} \frac{\langle \tilde{f}(S)u', S\tilde{f}(S)u' \rangle}{\langle \tilde{f}(S)u', \tilde{f}(S)u' \rangle} \end{aligned}$$

Notice that since U is upper triangular, the map taking $A \subset V_r$ to $A' = UA$ is a bijection among subspaces of V_r of given dimension. Since S and $\tilde{f}(S)$ are symmetric and commute,

$$\mu_{r,j}(F_{\tilde{f}}(S)) = \max_A \min_u \frac{\langle u, Sg(S)u \rangle}{\langle u, g(S)u \rangle},$$

where $g(x) = (\tilde{f}(x))^2$. The claim now follows from the inequality

$$\langle u, u \rangle \langle u, Sg(S)u \rangle - \langle u, Su \rangle \langle u, g(S)u \rangle \geq 0.$$

Diagonalize $S = Q^* M Q$ and $g(S) = Q^* g(\Lambda) Q$ and write $Qu = (x_1, \dots, x_n)$ so that

$$2(\langle u, u \rangle \langle u, Sg(S)u \rangle - \langle u, Su \rangle \langle u, g(S)u \rangle) = \sum_{k,\ell} (\mu_k - \mu_\ell)(g(\mu_k) - g(\mu_\ell)) x_k^2 x_\ell^2 \geq 0.$$

Consider now equality for the case $r = 1$. Notice that, by hypothesis, if $k \neq \ell$ then $(\mu_k - \mu_\ell)(g(\mu_k) - g(\mu_\ell)) > 0$. In the max-min formula for $\text{tr}(I_1 S) = \mu_{1,1}(S)$, it suffices to take $u = e_1$. Equality therefore holds only if Qe_1 is a canonical vector, which implies $(S)_{1,j} = 0$ for all $j > 1$. ■

Proof of Proposition 6.2: For all $s \in \mathcal{I}_i$ and any distinct eigenvalues λ_j and λ_k , $|\lambda_j - \lambda_i| < |\lambda_k - \lambda_i|$ if and only if $L_i(\lambda_j) < L_i(\lambda_k)$. For $s \in \mathcal{I}_i$, $f(x) = x - s$, $h(x) = L_i(x)$ and $\mu_j = L_i(\lambda_j)$, define $\tilde{f} : \mathbb{R} \rightarrow \mathbb{R}$ as in Lemma 6.3. The function \tilde{f} satisfies the hypothesis of Lemma 6.4: $\mu_j < \mu_k$ implies $|\tilde{f}(\mu_j)| < |\tilde{f}(\mu_k)|$. Thus, by

Lemma 6.4, $\text{tr}(WF_f(S)) \geq \text{tr}(WS)$ for all $S \in \mathcal{O}_\Sigma$. For $T \in \mathcal{D}_{\Lambda, \epsilon_{\text{ap}}}^i$, take $S = h(T)$: by Lemma 6.3, $F_{\bar{f}}(h(T)) = h(F_f(T))$ and therefore $\text{tr}(Wh(F_f(T))) \geq \text{tr}(Wh(T))$. Again by Lemma 6.4, equality happens only if T is diagonal. Thus, H_i is a height function. Finally, choosing δ_H sufficiently small guarantees that H_i is large in $\mathcal{D}_{\Lambda, 0}^i$ and small in $\partial\mathcal{D}_{\Lambda, \epsilon_{\text{ap}}}^i$, completing the proof. \blacksquare

Thus, simple shift strategies admit height functions near the deflation set. Our reason for constructing a height function is to control the time the sequence $(F_\sigma^k(T))$ stays in a compact set.

Assuming Λ to be a.p. free, for a shift strategy $\sigma : \mathcal{T}_\Lambda \rightarrow \mathbb{R}$ set $\epsilon_\sigma = \epsilon_{\text{ap}}/(1 + C_\sigma)$ (where C_σ is the constant in the definition of a simple shift strategy). Notice that $T \in \mathcal{D}_{\Lambda, \epsilon_\sigma}^i$ implies $\sigma(T) \in \mathcal{I}_i = [\lambda_i - \epsilon_{\text{ap}}, \lambda_i + \epsilon_{\text{ap}}]$.

Corollary 6.5 *Let Λ be a real diagonal $n \times n$ a.p. free matrix, σ a simple shift strategy and $\mathcal{D}_{\Lambda, \epsilon_\sigma}^i$ as above. Let $\mathcal{K} \subset \mathcal{D}_{\Lambda, \epsilon_\sigma}^i$ be a compact set with no diagonal matrices: there exists $K \in \mathbb{N}$ such that for all $T \in \mathcal{D}_{\Lambda, \epsilon_\sigma}^i$ there are at most K points of the form $F_\sigma^k(T)$ in \mathcal{K} .*

The plan is to take \mathcal{K} containing $\mathcal{S}_\sigma \cap \mathcal{D}_{\Lambda, \epsilon_\sigma}^i$: the hypothesis in Theorem 3 that diagonal matrices do not belong to the singular support \mathcal{S}_σ is then natural.

Proof: Let m_- be the minimum jump in \mathcal{K} and m_+ the size of the image of H_i :

$$m_- = \inf_{T \in \mathcal{K}, s \in \mathcal{I}_i} H_i(F_s(T)) - H_i(T), \quad m_+ = \sup_{T \in \mathcal{D}_{\Lambda, \epsilon_\sigma}^i} H_i(T) - \inf_{T \in \mathcal{D}_{\Lambda, \epsilon_\sigma}^i} H_i(T).$$

By Proposition 6.2 and the compactness of $\mathcal{K} \times \mathcal{I}_i$, $s > 0$: take K such that $Km_- > m_+$. For a given T , let $X = \{k \in \mathbb{N} \mid F_\sigma^k(T) \in \mathcal{K}\}$: we have

$$m_+ \geq \sum_{k \in X} H_i(F_\sigma^{k+1}(T)) - H_i(F_\sigma^k(T)) \geq |X|m_-$$

and therefore $|X| < K$. \blacksquare

Proof of Theorem 3: Let $\mathcal{K}_1, \mathcal{K}_2 \subset \mathcal{D}_{\Lambda, \epsilon_\sigma}^i$ be compact sets with $\mathcal{K}_1 \cup \mathcal{K}_2 = \mathcal{D}_{\Lambda, \epsilon_\sigma}^i$, $\mathcal{S}_\sigma \cap \mathcal{D}_{\Lambda, 0}^i$ disjoint from \mathcal{K}_1 and with no diagonal matrices in \mathcal{K}_2 . By Theorem 2, there exists $C_{\mathcal{K}_1} > 0$ such that $|b(F_\sigma(T))| \leq C_{\mathcal{K}_1}|b(T)|^3$ for all $T \in \mathcal{K}_1$. By Corollary 6.5, there exists $K_2 \in \mathbb{N}$ such that, given $T \in \mathcal{D}_{\Lambda, \epsilon_\sigma}^i$, at most K_2 points of the form $F_\sigma^k(T)$ belong to \mathcal{K}_2 . In particular, there are at most K_2 values of k for which the estimate $|b(F_\sigma^{k+1}(T))| \leq C_{\mathcal{K}_1}|b(F_\sigma^k(T))|^3$ does not hold. \blacksquare

7 Convergence rates for a.p. spectra

The aim of this section is to prove Theorem 4. An a.p. matrix $T \in \mathcal{T}$ with simple spectrum is *strong a.p.* if three consecutive eigenvalues are in arithmetic progression and *weak a.p.* otherwise.

In the a.p. free case discussed in the previous sections, for an initial condition $T \in \mathcal{D}_{\Lambda, \epsilon}^i$, the sequence $F_\sigma^k(T)$ converges to a diagonal matrix; this follows from the fact that $\sigma(T) \approx \lambda_i$ for $T \in \mathcal{D}_{\Lambda, \epsilon}^i$. For weak a.p. spectra, convergence to a diagonal matrix may not occur.

Assume Λ to be weak a.p. Let $b_2(T) = T_{n-1, n-2}$ be the second-last subdiagonal entry; for consistency, write $b_1(T) = b(T)$. For any i , there exists a unique index $c(i)$ such that $\lambda_{c(i)}$ is the eigenvalue closest to λ_i . As we shall see, if $T \in \mathcal{D}_{\Lambda, \epsilon}^i$ then

$$\lim_{k \rightarrow \infty} b_1(F_\sigma^k(T)) = \lim_{k \rightarrow \infty} b_2(F_\sigma^k(T)) = 0, \quad \lim_{k \rightarrow \infty} (F_\sigma^k(T))_{n, n} = \lambda_i;$$

furthermore, if T is unreduced then

$$\lim_{k \rightarrow \infty} (F_\sigma^k(T))_{n-1, n-1} = \lambda_{c(i)}.$$

We begin with a technical lemma concerning the dynamics of steps F_s . Item (b) is a variation of the power method argument used to study the convergence of lower entries under QR steps.

Lemma 7.1 *Let $M = \text{diag}(\mu_1, \dots, \mu_m)$ be a real diagonal matrix with simple spectrum and $\mathcal{T}_M \subset \mathcal{T}$ be the manifold of real $m \times m$ tridiagonal matrices similar to M . Let $I \subset \mathbb{R}$ be a compact interval. Assume that there exists j , $1 \leq j \leq m$, such that*

$$\mu_j \notin I, \quad \max_{s \in I} |\mu_j - s| < \min_{k \neq j, s \in I} |\mu_k - s|.$$

Let $\mathcal{D}_{M, \epsilon}^j \subset \mathcal{T}_M$ be the j -th deflation neighborhood.

- (a) *There exist $\epsilon > 0$ and $C \in (0, 1)$ such that for all $\epsilon' \in (0, \epsilon)$ and $s \in I$ we have $F_s(\mathcal{D}_{M, \epsilon'}^j) \subset \mathcal{D}_{M, C\epsilon'}^j$.*
- (b) *Consider $T_0 \in \mathcal{T}_M$ unreduced, a sequence (s_k) of elements of I and $\epsilon > 0$. Define $T_{k+1} = F_{s_k}(T_k)$. Then there exists k such that $T_k \in \mathcal{D}_{M, \epsilon}^j$.*

This will be used to study $b_2(T)$ for $T \in \mathcal{D}_{\Lambda, \epsilon}^i$, setting $I = [\lambda_i - \epsilon, \lambda_i + \epsilon]$, $j = c(i)$, $M = \Lambda_i = \text{diag}(\lambda_1, \dots, \lambda_{i-1}, \lambda_{i+1}, \dots, \lambda_n)$, with the natural identification between \mathcal{T}_M and $\mathcal{D}_{\Lambda, 0}^i$.

Proof: Let $\tilde{C} \in (0, 1)$ be such that

$$\max_{s \in I} |\mu_j - s| < \tilde{C} \min_{k \neq j, s \in I} |\mu_k - s|.$$

Write

$$r(s, T) = \frac{(R_\star)_{m, m}}{(R_\star)_{m-1, m-1}}, \quad T - sI = Q_\star R_\star.$$

Recall from Lemma 2.2 and Corollary 4.3 that $b(F_s(T)) = r(s, T) b(T)$. We claim that for all $T \in \mathcal{D}_{M, 0}^j$ and $s \in I$, $|r(s, T)| \leq \tilde{C}$. Since $T \in \mathcal{D}_{M, 0}^j$, $|(R_\star)_{m, m}| = |\mu_j - s|$. Let R_- be the leading principal minor of R_\star of order $m - 1$: its singular values are $|\mu_k - s|$, $k \neq j$. In particular, all singular values are larger than $|(R_\star)_{m, m}|/\tilde{C}$. Thus

$$|(R_\star)_{m-1, m-1}| = \|e_{m-1}^* R_-\| \geq \frac{|(R_\star)_{m, m}|}{\tilde{C}} \|e_{m-1}\| = \frac{|(R_\star)_{m, m}|}{\tilde{C}},$$

proving our claim. Take $C = (1 + \tilde{C})/2$: by continuity, for sufficiently small $\epsilon > 0$, we have $|r(s, T)| < C$ for all $T \in \mathcal{D}_{M, \epsilon}^j$, $s \in I$. Thus, for $T \in \mathcal{D}_{M, \epsilon}^j$ and $s \in I$, $|b(F_s(T))| \leq C |b(T)|$; item (a) follows.

For item (b), write $T_{k+1} = Q_k^* T_k Q_k$ where $T_k - s_k I = Q_k R_k$ is a $Q_\star R_\star$ decomposition. Notice that, by hypothesis, I is disjoint from the spectrum so that $T_0 - s_0 I$ is invertible. We have $(T_0 - s_0 I)^{-1} = R^{-1} Q_0^*$ so the rows of Q_0^* are obtained from those of $(T_0 - s_0 I)^{-1}$ by Gram-Schmidt from bottom to top. In particular, $Q_0 e_m = c_0 (T_0 - s_0 I)^{-1} e_m$, $c_0 > 0$. More generally, we claim that

$$P_k e_m = c(T_0 - s_{k-1} I)^{-1} \cdots (T_0 - s_1 I)^{-1} (T_0 - s_0 I)^{-1} e_m, \\ c > 0, \quad P_k = Q_0 Q_1 \cdots Q_{k-1} \in SO(m).$$

Indeed, by induction and using that $T_1 = Q_0^* T_0 Q_0$,

$$P_k e_m = c' Q_0 (T_1 - s_{k-1} I)^{-1} \cdots (T_1 - s_1 I)^{-1} e_m \\ = c' (T_0 - s_{k-1} I)^{-1} \cdots (T_0 - s_1 I)^{-1} Q_0 e_m \\ = c(T_0 - s_{k-1} I)^{-1} \cdots (T_0 - s_1 I)^{-1} (T_0 - s_0 I)^{-1} e_m.$$

(Integrability of the Toda lattice is present here yet another time.) For $\alpha = 1, \dots, m$, let v_α be the unit eigenvector associated to μ_α . We claim that

$$\lim_{k \rightarrow \infty} P_k e_m = \pm v_j.$$

Indeed, write $e_m = \sum_{\alpha=1}^m a_\alpha v_\alpha$, where $a_\alpha = \langle v_\alpha, e_m \rangle$ is the last coordinate of v_α . It is well known that the last coordinates of the eigenvectors v_α of the unreduced matrix T are nonzero: in particular, $a_j \neq 0$; assume without loss $a_j > 0$. We have

$$\begin{aligned} P_k e_m &= c(T_0 - s_{k-1}I)^{-1} \cdots (T_0 - s_1I)^{-1} (T_0 - s_0I)^{-1} e_m \\ &= c \sum_{\alpha=1}^m \frac{a_\alpha}{(\mu_\alpha - s_{k-1}) \cdots (\mu_\alpha - s_0)} v_\alpha = c_k \left(v_j + \sum_{\alpha \neq j} b_{k,\alpha} v_\alpha \right), \\ c_k &> 0, \quad b_{k,\alpha} = \frac{a_\alpha}{a_j} \frac{\mu_j - s_{k-1}}{\mu_\alpha - s_{k-1}} \cdots \frac{\mu_j - s_0}{\mu_\alpha - s_0}. \end{aligned}$$

Since $|\mu_j - s_{k-1}|/|\mu_\alpha - s_{k-1}| < \tilde{C}$ we have $|b_{k,\alpha}| \leq (\tilde{C})^k |a_\alpha/a_j|$ and therefore $\lim_{k \rightarrow \infty} b_{k,\alpha} = 0$, proving the claim. We have

$$\begin{aligned} \lim_{k \rightarrow \infty} b(T_k) &= \lim_{k \rightarrow \infty} (T_k)_{m,m-1} = \lim_{k \rightarrow \infty} e_{n-1}^* T_k e_m = \lim_{k \rightarrow \infty} (P_k e_{m-1})^* T_0 (P_k e_m) = \\ &= \lim_{k \rightarrow \infty} (P_k e_{m-1})^* \mu_j (P_k e_m) + \lim_{k \rightarrow \infty} (P_k e_{m-1})^* (T_0 - \mu_j I) (P_k e_m). \end{aligned}$$

The first limit in the last expression is zero because $P_k e_{m-1} \perp P_k e_m$; the second is zero because $P_k e_{m-1}$ is bounded and

$$\lim_{k \rightarrow \infty} (T_0 - \mu_j I) (P_k e_m) = (T_0 - \mu_j I) \lim_{k \rightarrow \infty} (P_k e_m) = (T_0 - \mu_j I) v_j = 0.$$

■

Consider the *double deflation set* $\mathcal{C}_{\Lambda,0} \subset \mathcal{D}_{\Lambda,0} \subset \mathcal{T}_\Lambda$:

$$\mathcal{C}_{\Lambda,0} = \{T \in \mathcal{T}_\Lambda \mid b_1(T) = b_2(T) = 0\}.$$

For Wilkinson's strategy ω , it turns out that the set $\mathcal{C}_{\Lambda,0}$ is disjoint from the singular support \mathcal{S}_ω . More generally, if a shift strategy σ satisfies $\mathcal{C}_{\Lambda,0} \cap \mathcal{S}_\sigma = \emptyset$ then cubic convergence of F_σ holds even for weak a.p. spectra: this is Theorem 4, which we prove below.

In [9], we show examples of unreduced tridiagonal 3×3 matrices with spectrum $-1, 0, 1$ for which Wilkinson's shift F_ω converges quadratically to a reduced but not diagonal matrix in the singular support \mathcal{S}_ω . Similarly, we conjecture that for strong a.p. diagonal $n \times n$ matrices Λ there exists a set $\mathcal{X} \subset \mathcal{T}_\Lambda$ of Hausdorff codimension 1 of unreduced matrices T for which $F_\omega^k(T)$ converges quadratically to a matrix in $\mathcal{S}_\omega \cap \mathcal{D}_{\Lambda,0}$ with $T_{n-1,n-2} \neq 0$.

With the natural identification between $\mathcal{D}_{\Lambda,0}^i$ and \mathcal{T}_{Λ_i} , we may consider $\mathcal{D}_{\Lambda_i,\epsilon_2}^j$ to be a subset of $\mathcal{D}_{\Lambda,0}^i$. Let

$$\mathcal{C}_{\Lambda,\epsilon_2,\epsilon_1}^{j,i} = \mathcal{D}_{\Lambda,\epsilon_1}^i \cap \Pi_i^{-1}(\mathcal{D}_{\Lambda_i,\epsilon_2}^j).$$

For small $\epsilon_1, \epsilon_2 > 0$, $T \in \mathcal{C}_{\Lambda,\epsilon_2,\epsilon_1}^{j,i}$ implies

$$T_{n-1,n-1} \approx \lambda_j, \quad T_{n,n} \approx \lambda_i, \quad b_1(T) \leq \epsilon_1, \quad b_2(T) \approx 0.$$

These compact sets turn out to be manifolds with corners but we shall neither prove nor use this fact. Lemma 7.1 can be rephrased in terms of the sets $\mathcal{C}_{\Lambda,\epsilon_2,\epsilon_1}^{j,i}$.

Corollary 7.2 *Let Λ to be weak a.p. spectrum and σ be a simple shift strategy. There exists $\epsilon > 0$ such that, for all i and for all $\epsilon_1 \in (0, \epsilon)$:*

- (a) *there exists $C \in (0, 1)$ such that, for all sufficiently small $\epsilon_2 > 0$ we have $F_\sigma(\mathcal{C}_{\Lambda, \epsilon_2, \epsilon_1}^{c(i), i}) \subset \mathcal{C}_{\Lambda, C\epsilon_2, \epsilon_1}^{c(i), i}$;*
- (b) *for all unreduced $T \in \mathcal{D}_{\Lambda, \epsilon}^i$ and for all $\epsilon_1, \epsilon_2 > 0$ there exists k such that $F_\sigma^k(T) \in \mathcal{C}_{\Lambda, \epsilon_2, \epsilon_1}^{c(i), i}$.*

Proof: Combine Lemma 7.1 with $\Pi_i \circ F_s = F_s \circ \Pi_i$ (Proposition 4.1). ■

Proof of Theorem 4: From the hypothesis that $\mathcal{C}_{\Lambda, 0}$ and \mathcal{S}_σ are disjoint it follows that, for sufficiently small $\epsilon_1, \epsilon_2 > 0$, the shift strategy σ is smooth in $\mathcal{C}_{\Lambda, \epsilon_2, \epsilon_1}^{c(i), i}$. As in Lemma 5.2, from a Taylor expansion around $T_0 \in \mathcal{D}_{\Lambda, 0}^i$, there exists C_2 such that $|\sigma(T)| \leq C_2|b_1(T)|^2$ for all $T \in \mathcal{C}_{\Lambda, \epsilon_2, \epsilon_1}^{c(i), i}$. As in the proof of Theorem 2, there exists C_3 such that $|b_1(F_\sigma(T))| \leq C_3|b_1(T)|^3$ for all $T \in \mathcal{C}_{\Lambda, \epsilon_2, \epsilon_1}^{c(i), i}$. From item (a) of Corollary 7.2, $\mathcal{C}_{\Lambda, \epsilon_2, \epsilon_1}^{c(i), i}$ is invariant under F_σ ; from item (b), for all unreduced $T \in \mathcal{D}_{\Lambda, \epsilon}^i$ (where ϵ is sufficiently small) there exists K such that, for all $k > K$, $F_\sigma^k(T) \in \mathcal{C}_{\Lambda, \epsilon_2, \epsilon_1}^{c(i), i}$, completing the proof. ■

8 Two counterexamples

In this section we present two examples which show that natural strengthenings of Theorems 3 and 4 do not hold for Wilkinson's strategy ω .

We use the notation of Section 3. In Figure 4, where $\Lambda = \text{diag}(1, 2, 4)$, we indicate a sequence $\tilde{F}_\omega^k(T)$ which enters the deflation neighborhood $\mathcal{D}_{\Lambda, \epsilon}^i$ near one diagonal matrix but travels within the neighborhood towards another diagonal matrix. Theorem 2 guarantees the cubic decay of the $(3, 2)$ entry whenever $\tilde{F}_\omega^k(T)$ stays away from the singular support \mathcal{S}_ω . Consistently with Theorem 3, this happens for practically all values of k . Notice however that no uniform bound exists on the number of iterations needed to reach (a neighborhood of) \mathcal{S}_ω . As proved in [9], in this instance cubic decay does not hold. More precisely, it is *not* true that given an a.p. free matrix Λ there exist $C > 0$ and K such that $|b(F_\omega^{k+1}(T))| \leq C|b(F_\omega^k(T))|^3$ for all $k > K$.




Figure 4: We may have $F_\omega^k(T) \in \mathcal{S}_\omega$ for large values of k .

Consider now the weak a.p. spectrum $\Lambda = \text{diag}(-1, 0, 0.3, 1)$ and

$$T_0 = \begin{pmatrix} 0.3 & 0 \\ 0 & S_0 \end{pmatrix} \in \mathcal{T}_\Lambda$$

where $S_0 \in \mathcal{T}_{\Lambda_3}$, $\Lambda_3 = \text{diag}(-1, 0, 1)$, is an example of unreduced matrix obtained in [9] for which convergence is strictly quadratic, i.e.,

$$C_-|b(F_\omega^k(S_0))|^2 < |b(F_\omega^{k+1}(S_0))| < C_+|b(F_\omega^k(S_0))|^2,$$

for all k , where $0 < C_- < C_+$. Trivially, the analogous estimate holds for $b(F_\omega^k(T_0))$. By sheer continuity, given K , there exists $\epsilon > 0$ such that if $T \in \mathcal{T}_\Lambda$ satisfies $\|T - T_0\| < \epsilon$ then

$$C_- |b(F_\omega^k(T))|^2 < |b(F_\omega^{k+1}(T))| < C_+ |b(F_\omega^k(T))|^2$$

still holds for all $k < K$. Thus, the uniform estimate in Theorem 3 fails for weak a.p. spectra, even for unreduced matrices.

9 Appendix: Proof of Theorem 1

Recall from Section 2 that $E_j \in \mathcal{E}$ is defined by

$$(E_j)_{i,i} = \begin{cases} 1, & i \leq j, \\ -1, & i > j; \end{cases}$$

the involutions η_j are defined by $\eta_j(T) = E_j T E_j$, which differs from T only in the sign of the j -th subdiagonal coordinate. Let $\mathcal{M}_j \subset \mathcal{T}_\Lambda$ be the *mirror*, i.e., the set of fixed points of η_j : for $T \in \mathcal{T}_\Lambda$ we have $T \in \mathcal{M}_j$ if and only if $(T)_{j+1,j} = 0$. Let S_n be the symmetric group of permutations π of the set $\{1, 2, \dots, n\}$. For $\pi \in S_n$, let $\mathcal{M}_{j,\pi} \subset \mathcal{M}_j$ be the set of matrices for which the eigenvalues of the top $j \times j$ principal subblock are $\lambda_{\pi(1)}, \dots, \lambda_{\pi(j)}$ so that

$$\mathcal{M}_{j,\pi} \approx \mathcal{T}_{\text{diag}(\lambda_{\pi(1)}, \dots, \lambda_{\pi(j)})} \times \mathcal{T}_{\text{diag}(\lambda_{\pi(j+1)}, \dots, \lambda_{\pi(n)})}.$$

Thus, \mathcal{M}_j is a submanifold of codimension 1 with $\binom{n}{j}$ connected components $\mathcal{M}_{j,\pi}$. The diagonal matrices in \mathcal{T}_Λ are labeled by $\pi \in S_n$: let

$$\Lambda^\pi = \text{diag}(\lambda_{\pi(1)}, \dots, \lambda_{\pi(n)}).$$

Let $\tilde{\mathcal{J}}_\Lambda \subset \mathcal{T}_\Lambda$ be the set of tridiagonal matrices with nonnegative subdiagonal entries. The set $\tilde{\mathcal{J}}_\Lambda$ is homeomorphic to the *permutohedron* \mathcal{P}_Λ ([16]), the convex hull of the points

$$v_\pi = (\lambda_{\pi^{-1}(1)}, \dots, \lambda_{\pi^{-1}(n)}) \in \mathbb{R}^n, \quad \pi \in S_n;$$

the vertices of \mathcal{P}_Λ are v_π . An explicit homeomorphism takes $T = Q^* \Lambda Q$ to the vector in \mathbb{R}^n whose j -th coordinate is $(Q \Lambda Q^*)_{jj}$; this map takes Λ^π to v_π ([2]). We use this map to endow $\tilde{\mathcal{J}}_\Lambda$ with a combinatorial structure of vertices, faces and hyperfaces: in particular, vertices of $\tilde{\mathcal{J}}_\Lambda$ are diagonal matrices. It turns out that the hyperfaces of $\tilde{\mathcal{J}}_\Lambda$ are the intersections $\mathcal{M}_{j,\pi} \cap \tilde{\mathcal{J}}_\Lambda$.

Lemma 9.1 *Let $\mathcal{P} \subset \mathbb{R}^n$ be the convex hull of a finite set. Let $\tilde{F} : \mathcal{P} \rightarrow \mathcal{P}$ be a continuous function. Assume that for any hyperface $\mathcal{Q} \subset \mathcal{P}$ we have $\tilde{F}(\mathcal{Q}) \subset \mathcal{Q}$. Then \tilde{F} is surjective.*

Proof: The dimension of a convex subset of \mathbb{R}^n is the dimension of the affine subspace spanned by its vertices. Notice that any face (of any dimension) is the intersection of hyperfaces and therefore also invariant under \tilde{F} .

We use relative homology: if the dimension of \mathcal{P} is d then $H_d(\mathcal{P}, \partial\mathcal{P}) = \mathbb{Z}$; we prove that $\tilde{F}_* : H_d(\mathcal{P}, \partial\mathcal{P}) \rightarrow H_d(\mathcal{P}, \partial\mathcal{P})$ is the identity. This implies the lemma: if x_0 is an interior point of \mathcal{P} not in the image of \tilde{F} then since $H_d(\mathcal{P}, \mathcal{P} \setminus \{x_0\}) = H_d(\mathcal{P}, \partial\mathcal{P})$ we have $\tilde{F}_* = 0$, a contradiction.

The proof of the claim is by induction on the dimension d of \mathcal{P} . The case $d = 0$ is trivial; in the case $d = 1$ the polytope \mathcal{P} is an interval and \tilde{F} takes each endpoint

to itself and again the claim is easy. In general, let Q be a hyperface of \mathcal{P} so that the dimension of Q is $d - 1$ and, by induction, $\tilde{F}_* : H_{d-1}(Q, \partial Q) \rightarrow H_{d-1}(Q, \partial Q)$ is the identity. We have $Q \subset \partial\mathcal{P}$ and $H_{d-1}(Q, \partial Q) = H_{d-1}(\partial\mathcal{P}, \partial Q \cup (\partial\mathcal{P} \setminus Q)) = H_{d-1}(\partial\mathcal{P})$ and therefore $\tilde{F}_* : H_{d-1}(\partial\mathcal{P}) \rightarrow H_{d-1}(\partial\mathcal{P})$ is the identity. Since \mathcal{P} is contractible, the long exact sequence for relative homology implies that $\tilde{F}_* : H_d(\mathcal{P}, \partial\mathcal{P}) \rightarrow H_d(\mathcal{P}, \partial\mathcal{P})$ is the identity, completing the proof. ■

Proof of Theorem 1: For (a), first notice that the condition $F \circ \eta_i = \eta_i \circ F$ implies $F(\mathcal{M}_i) \subseteq \mathcal{M}_i$. Since diagonal matrices are fixed points this implies $F(\mathcal{M}_{i,g}) \subseteq \mathcal{M}_{i,g}$. Restrict F to $\tilde{\mathcal{J}}_\Lambda$ and drop signs to define a continuous map $\tilde{F} : \tilde{\mathcal{J}}_\Lambda \rightarrow \tilde{\mathcal{J}}_\Lambda$ which keeps each hyperface of $\tilde{\mathcal{J}}_\Lambda$ invariant. By Lemma 9.1, \tilde{F} is surjective and therefore (by equivariance) so is F .

For (b), let $\mathcal{B}^i \subset \mathcal{T}_\Lambda$ be the basins of attraction of each invariant neighborhood $\text{int}(\mathcal{K}_i)$, i.e., $T \in \mathcal{B}^i$ if there exists $k \in \mathbb{N}$ such that $F_\sigma^k(T) \in \text{int}(\mathcal{K}_i)$. The sets \mathcal{B}^i are clearly disjoint with $\mathcal{K}_i \subset \mathcal{B}^i$. They are also open subsets of \mathcal{T}_Λ since $\mathcal{B}^i = \bigcup_k F^{-k}(\text{int}(\mathcal{K}_i))$. Since \mathcal{T}_Λ is connected there exists $T \notin \bigcup_i \mathcal{B}^i$ and we are done. ■

References

- [1] Bloch, A. M., Brockett, R. W. and Ratiu, T., Completely integrable gradient flows, *Comm. Math. Phys.*, 147, 57-74, 1992.
- [2] Bloch, A. M., Flaschka, H. and Ratiu, T., A convexity theorem for isospectral manifolds of Jacobi matrices in a compact Lie algebra, *Duke Math. J.*, 61, 41-65, 1990.
- [3] Deift, P., Nanda, T., Tomei, C., Differential equations for the symmetric eigenvalue problem, *SIAM J. Num. Anal.* 20, 1-22, 1983.
- [4] Deift, P., Rivera, S., Tomei, C., Watkins, D., A monotonicity property for Toda-type flows, *SIAM J. Matrix Anal. Appl.*, 12, 463-468, 1991.
- [5] Demmel, J. W., *Applied Numerical Linear Algebra*, SIAM, Philadelphia, 1997.
- [6] Flaschka, H., The Toda lattice, I, *Phys. Rev. B* 9, 1924-1925, 1974.
- [7] Hoffmann, W. and Parlett, B., A new proof of global convergence for the tridiagonal QL algorithm, *SIAM J. Num. Anal.* 15 (1978), 929-937.
- [8] Leite, R. S., Saldanha, N.C. and Tomei, C., An atlas for tridiagonal isospectral manifolds, *Lin. Alg. Appl.* 429 (2008), 387-402.
- [9] Leite, R. S., Saldanha, N.C. and Tomei, C., The Asymptotics of Wilkinson's shift: Loss of Cubic Convergence, *FoCM* 10 (1), 15-36, 2010. doi:10.1007/s10208-009-9047-3
- [10] Leite, R. S., Saldanha, N.C. and Tomei, C., The Asymptotics of Wilkinson's shift iteration, arXiv:math/0412493v2.
- [11] Leite, R. S. and Tomei, C., Parametrization by polytopes of intersections of orbits by conjugation, *Lin. Alg. and Appl.* 361, 223-246, 2003.
- [12] Moerbeke, P. van, The spectrum of Jacobi matrices, *Inventiones Math.*, 37, 45-81, 1976.

- [13] Moser, J., Finitely many mass points on the line under the influence of an exponential potential, In: Dynamic systems theory and applications, (ed. J. Moser) 467-497, New York, 1975.
- [14] Parlett, B. N., The Symmetric Eigenvalue Problem, Classics in Applied Mathematics 20, SIAM, 1997.
- [15] Sternberg, S., Local contractions and a theorem of Poincaré, Amer. J. Math., 79, 809-824, 1957.
- [16] Tomei, C., The Topology of Manifolds of Isospectral Tridiagonal Matrices, Duke Math. J., 51, 981-996, 1984.
- [17] Wilkinson, J. H., The algebraic eigenvalue problem, Oxford University Press, 1965.

Ricardo S. Leite, Departamento de Matemática, UFES
Av. Fernando Ferrari, 514, Vitória, ES 29075-910, Brazil

Nicolau C. Saldanha and Carlos Tomei, Departamento de Matemática, PUC-Rio
R. Marquês de S. Vicente 225, Rio de Janeiro, RJ 22453-900, Brazil

rsleite@pq.cnpq.br

saldanha@puc-rio.br; <http://www.mat.puc-rio.br/~nicolau/>

tomei@mat.puc-rio.br